



# **STIC Search Report**

## **Biotech-Chem Library**

**STIC Database Tracking Number: 94933**

**TO: Josephine Young**  
**Location: 8d04 / 8b19**  
**Wednesday, June 04, 2003**  
**Art Unit: 1623**  
**Phone: 605-1201**  
**Serial Number: 09 / 844450**

**From: Jan Delaval**  
**Location: Biotech-Chem Library**  
**CM1-1E07**  
**Phone: 308-4498**  
**jan.delaval@uspto.gov**

### **Search Notes**

Jan Delaval  
Reference Librarian  
Biotechnology & Chemical Library  
CM1 1E07 - 703-308-4498  
jan.delaval@uspto.gov



=&gt; d his

(FILE 'HOME' ENTERED AT 08:06:33 ON 04 JUN 2003)  
SET COST OFF

FILE 'HCAPLUS' ENTERED AT 08:06:47 ON 04 JUN 2003

Jan Delaval  
Reference Librarian  
Biotechnology & Chemical Library  
CM1 1E07 - 703-308-4498  
jan.delaval@uspto.gov

E US20020028786/PN  
L1 1 S E3  
E WO2001-US13931/AP, PRN  
L2 1 S E3, E4  
E US2000-233263/AP, PRN  
L3 1 S E5  
E US2000-233025/AP, PRN  
L4 1 S E5  
E US2000-230263/AP, PRN  
L5 1 S E5  
L6 1 S L1-L5  
E HEALTHPARTNER/PA, CS  
L7 12 S E5-E11  
E HEALTH PARTNER/PA, CS  
L8 23 S (HEALTH (L) PARTN?) /PA, CS  
SEL DN AN 23  
L9 1 S L8 AND E1-E3  
L10 13 S L7, L9  
E FREY W/AU  
L11 167 S E3, E7-E9, E25, E29-E33  
E FAWCETT J/AU  
L12 33 S E3, E13  
E FAWCETT JOHN/AU  
L13 171 S E3, E6, E7  
L14 48821 S ?PYROPHOS?  
L15 144 S ?PYRO PHOS?  
E PYROPHOS/CT  
E E16+ALL  
L16 851 S E1  
E E2+ALL  
L17 8258 S E10+NT  
E E8+ALL  
L18 24848 S E4+NT  
E E3+ALL  
L19 219974 S E3+NT  
L20 2 S L10 AND L14-L19  
L21 3 S L11-L13 AND L14-L19  
L22 3 S L6, L20, L21  
L23 2 S L22 NOT RHODIUM/TI  
E ALZHEIMER/CT  
L24 11589 S E3-E20  
E E9+ALL  
L25 11603 S E6, E5+NT  
L26 11213 S E23+NT OR E24+NT OR E25+NT OR E26+NT OR E27+NT OR E28+NT OR E  
E E23+ALL  
L27 1188 S E8  
E E6+ALL  
L28 72281 S E7, E6+NT  
E E22+ALL  
E E5+ALL  
E E25+ALL  
L29 10751 S E9+NT  
E E8+ALL  
L30 2080 S E6, E5+NT  
L31 28838 S E15+NT OR E13+NT  
E E10+ALL  
E E27+ALL

L32 449 S E8-E11  
 E E14+ALL  
 E E28+ALL  
 E E3+ALL  
 E E29+ALL  
 L33 961 S E3  
 L34 3 S L10 AND L24-L33  
 L35 14 S L11-L13 AND L24-L33  
 L36 14 S L23,L34,L35  
 L37 4 S L36 AND ?PHOSPH?  
 L38 2 S L37 AND L23  
 SEL RN

FILE 'REGISTRY' ENTERED AT 08:21:28 ON 04 JUN 2003

L39 66 S E1-E67  
 L40 14 S L39 AND P/ELS  
 L41 3 S L40 AND (C4H8CL3O4P OR C2H8O7P2 OR C3H11NO7P2)  
 L42 1 S L40 AND H3O3P  
 L43 2 S L41 AND P>=2  
 L44 10 S L40 NOT L41,L42  
 L45 12 S L43,L44  
 L46 151 S (C6H16O18P4 OR C6H15O15P3 OR C6H14O12P2)/MF AND 46.150.1/RID  
 L47 9 S 2466-09-3 OR 10380-08-2 OR 13813-62-2 OR 12395-97-0 OR 29444-  
 L48 1 S 7664-38-2  
 L49 10 S L47,L48  
 SEL RN  
 L50 16644 S E68-E77/CRN  
 L51 2775 S L50 AND PMS/CI  
 L52 232 S L51 AND HOMOPOLYMER  
 L53 117 S L52 AND NR>=1  
 L54 115 S L52 NOT L53  
 L55 16 S L54 AND (LI OR H3N OR NA)  
 L56 4 S L55 AND 2/NC  
 L57 8 S L55 AND 3/NC  
 L58 1 S L57 AND K  
 L59 5 S L56,L58  
 L60 2543 S L51 NOT L52-L59  
 L61 1751 S L60 AND NR>=1  
 L62 792 S L60 NOT L61  
 L63 363 S L62 NOT (C2H4O OR C3H6O)  
 L64 142 S L63 NOT (N OR S OR SI)/ELS  
 L65 2697 S L51 AND H3O4P  
 L66 5 S L65 AND 1/NC  
 L67 2 S L40 NOT L45  
 L68 1 S L67 AND H3O3P  
 L69 22 S L45,L49,L68  
 L70 170 S L46,L69  
 L71 10 S (O7P2 OR O13P4 OR O16P5 OR O19P6 OR O22P7 OR O25P8 OR L28P9 O  
 L72 180 S L70,L71  
 E HEXAMETAPHOSPHATE/CN  
 L73 1 S E3  
 L74 2 S E4,E5  
 E OPOP/ES  
 L75 1511 S (OPOP OR OPOPOP OR OPOPOP OR OPOPOP OR OPOPOP OR OPOPOP  
 L76 177 S L75 AND 1/NR AND 1/NC  
 L77 162 S L76 NOT (ETHOXY OR METHOXY OR PROPOXY)  
 L78 134 S L77 NOT (TIS OR AYS OR MNS)/CI  
 L79 132 S L78 NOT CCS/CI  
 L80 50 S L79 NOT (C OR F OR CL OR S OR N)/ELS  
 L81 41 S L80 NOT RPS/CI  
 L82 35 S L81 NOT ION  
 L83 9 S L82 AND (H2O6P2 OR H3O8P3 OR H4O12P4 OR H5O15P5 OR H6O18P6 OR  
 L84 6 S L83 NOT 3-4/P

L85 3 S L83 NOT L84  
 L86 1 S L85 NOT OPOPOP/ES  
 L87 7 S L84,L86  
 L88 2 S L75 AND H9O27P9  
 L89 268 S L75 AND H3O9P3  
 L90 1 S L89 AND 1/NC  
 L91 8 S L87,L90  
 L92 188 S L72,L91  
 E GLASS/CN  
 L93 1 S E56  
 L94 189 S L92,L93

FILE 'HCAPLUS' ENTERED AT 09:07:16 ON 04 JUN 2003  
 L95 81393 S L94  
 L96 5383 S INOSITOL(L) (DIPHOSPH? OR TRIPHOS? OR TETRAPHOS? OR PENTAPHOS?  
 L97 82822 S IMIDODIPHOS? OR GUANYLIMIDODIPHOS? OR ADENYLYLIMIDODIPHOS? OR  
 L98 111 S (ETIDRONIC OR PAMIDRONIC)()ACID

FILE 'REGISTRY' ENTERED AT 09:08:54 ON 04 JUN 2003

FILE 'HCAPLUS' ENTERED AT 09:11:24 ON 04 JUN 2003  
 L99 991 S AMIDRONIC ACID OR (GUANYL? OR ADENYL?)()IMIDODIPHOSPH?  
 L100 156796 S L95-L98,L99  
 L101 1483 S L24-L33 AND L100  
 L102 17 S L10-L13 AND L100  
 L103 2 S L101 AND L102  
 L104 2 S L38,L103  
 L105 15 S L102 NOT L104  
 L106 91 S L100 (L) THU/RL AND L101  
 L107 66 S L95 (L) THU/RL AND L106  
 L108 12 S L107 AND ?ALZHEIM?  
 L109 9 S L107 AND ?AMYLO?  
 L110 19 S L108,L109  
 L111 1277 S L101 AND (PD<=20000501 OR PRD<=20000501 OR AD<=20000501)  
 L112 63 S L111 AND L106  
 L113 10 S L110 AND L112  
 SEL DN AN 2 3 4 5 6  
 L114 5 S E1-E15 AND L113  
 L115 53 S L112 NOT L113  
 SEL DN AN 16 28 27 42  
 L116 4 S E16-E27 AND L115  
 L117 10 S L114,L116,L104  
 SEL HIT RN

FILE 'REGISTRY' ENTERED AT 09:25:37 ON 04 JUN 2003  
 L118 16 S E28-E43  
 L119 15 S L118 NOT UNSPECIFIED

=> fil reg

FILE 'REGISTRY' ENTERED AT 09:26:07 ON 04 JUN 2003  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2003 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
 provided by InfoChem.

STRUCTURE FILE UPDATES: 3 JUN 2003 HIGHEST RN 524916-37-8  
 DICTIONARY FILE UPDATES: 3 JUN 2003 HIGHEST RN 524916-37-8

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 6, 2003

Please note that search-term pricing does apply when

conducting SmartSELECT searches.


Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. See HELP PROPERTIES for more information. See STNote 27, Searching Properties in the CAS Registry File, for complete details:  
<http://www.cas.org/ONLINE/STN/STNOTES/stnotes27.pdf>

=> d ide can tot l119

L119 ANSWER 1 OF 15 REGISTRY COPYRIGHT 2003 ACS  
 RN 85166-31-0 REGISTRY  
 CN D-myo-Inositol, 1,4,5-tris(dihydrogen phosphate) (9CI) (CA INDEX NAME)  
 OTHER CA INDEX NAMES:  
 CN Inositol triphosphate (6CI, 7CI)  
 OTHER NAMES:  
 CN D-myo-Inositol 1,4,5-triphosphate  
 CN D-myo-Inositol 1,4,5-trisphosphate  
 CN Inositol trisphosphate  
 FS STEREOSEARCH  
 DR 146952-67-2, 346624-32-6  
 MF C6 H15 O15 P3  
 CI COM  
 LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, BEILSTEIN\*, BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CANCERLIT, CAOLD, CAPLUS, CASREACT, CEN, CHEMCATS, CIN, CSChem, EMBASE, MEDLINE, PROMT, TOXCENTER, USPATFULL  
 (\*File contains numerically searchable property data)

Absolute stereochemistry. Rotation (-).



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

953 REFERENCES IN FILE CA (1957 TO DATE)  
 24 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
 955 REFERENCES IN FILE CAPLUS (1957 TO DATE)  
 18 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

REFERENCE 1: 138:332173  
 REFERENCE 2: 138:332169  
 REFERENCE 3: 138:183487  
 REFERENCE 4: 138:181193  
 REFERENCE 5: 138:131350  
 REFERENCE 6: 138:117878  
 REFERENCE 7: 138:86984

REFERENCE 8: 138:69849

REFERENCE 9: 138:33547

REFERENCE 10: 138:22933

L119 ANSWER 2 OF 15 REGISTRY COPYRIGHT 2003 ACS

RN 40391-99-9 REGISTRY

CN Phosphonic acid, (3-amino-1-hydroxypropylidene)bis- (9CI) (CA INDEX NAME)

OTHER NAMES:

CN (.alpha.-Hydroxy-.gamma.-aminopropylidene)diphosphonic acid

CN (3-Amino-1-hydroxypropylidene)-1,1-bisphosphonate

CN 3-Amino-1-hydroxypropane-1,1-diphosphonic acid

CN 3-Amino-1-hydroxypropylidenediphosphonic acid

CN ADP

CN AHPrBP

CN Amidronic acid

CN Pamidronic acid

CN Propane-1-hydroxy-3-amino-1,1-diphosphonic acid

FS 3D CONCORD

MF C3 H11 N O7 P2


CI COM

LC STN Files: ADISINSIGHT, ADISNEWS, ANABSTR, BEILSTEIN\*, BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CANCERLIT, CAPLUS, CASREACT, CBNB, CHEMCATS, CHEMINFORMRX, CHEMLIST, CSCHEM, DDFU, DRUGNL, DRUGPAT, DRUGU, DRUGUPDATES, EMBASE, IFICDB, IFIPAT, IFIUDB, IPA, MEDLINE, MRCK\*, PHAR, PROMT, SPECINFO, SYNTHLINE, TOXCENTER, USAN, USPAT2, USPATFULL, VETU

(\*File contains numerically searchable property data)

Other Sources: EINECS\*\*, WHO

(\*\*Enter CHEMLIST File for up-to-date regulatory information)



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

640 REFERENCES IN FILE CA (1957 TO DATE)

31 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

646 REFERENCES IN FILE CAPLUS (1957 TO DATE)

REFERENCE 1: 138:352034

REFERENCE 2: 138:348231

REFERENCE 3: 138:331682

REFERENCE 4: 138:331345

REFERENCE 5: 138:326535

REFERENCE 6: 138:326440

REFERENCE 7: 138:297381

REFERENCE 8: 138:292738

REFERENCE 9: 138:281598

REFERENCE 10: 138:281077

L119 ANSWER 3 OF 15 REGISTRY COPYRIGHT 2003 ACS

RN **34273-04-6** REGISTRY

CN 5'-Guanylic acid, monoanhydride with imidodiphosphoric acid (8CI, 9CI)  
(CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Imidodiphosphoric acid, monoanhydride with 5'-guanylic acid (8CI)

OTHER NAMES:

CN .beta.,.gamma.-Imidoguanosine-5'-triphosphate

CN 5'-Guanylyl imidodiphosphate

CN 5'-Guanylyliminodiphosphonate

CN 5-Guanylylimidodiphosphate

CN Guanosine 5'-(.beta.,.gamma.-imido)triphosphate

CN Guanosine 5'-(.beta.,.gamma.-imino)triphosphate

CN Guanylyl imidodiphosphate

CN Guanylyl-.beta.,.gamma.-imidodiphosphate

CN Guanylyl-5'-(.beta.,.gamma.-imido)diphosphate

FS STEREOSEARCH


DR 94725-19-6, 104062-58-0, 104838-92-8, 92836-22-1, 110378-98-8

MF C10 H17 N6 O13 P3

CI COM

LC STN Files: AGRICOLA, BEILSTEIN\*, BIOBUSINESS, BIOSIS, BIOTECHNO, CA,  
CANCERLIT, CAPLUS, DDFU, DRUGU, EMBASE, MEDLINE, TOXCENTER, USPATFULL  
(\*File contains numerically searchable property data)

Absolute stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

2044 REFERENCES IN FILE CA (1957 TO DATE)

47 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

2045 REFERENCES IN FILE CAPLUS (1957 TO DATE)

REFERENCE 1: 138:316710

REFERENCE 2: 138:248790

REFERENCE 3: 138:163487

REFERENCE 4: 138:148027

REFERENCE 5: 138:117741

REFERENCE 6: 137:381427

REFERENCE 7: 137:380285

REFERENCE 8: 137:364731

REFERENCE 9: 137:197091

REFERENCE 10: 137:151462

L119 ANSWER 4 OF 15 REGISTRY COPYRIGHT 2003 ACS

RN 27590-04-1 REGISTRY

CN Imidodiphosphoric acid (6CI, 7CI, 8CI, 9CI) (CA INDEX NAME)

OTHER NAMES:

CN Imidobisphosphonic acid

CN Imidodiphosphonic acid

CN Phosphonic acid, iminobis-

FS 3D CONCORD

DR 86960-45-4

MF H5 N O6 P2

CI COM

LC STN Files: BIOSIS, CA, CAOLD, CAPLUS, CHEMLIST, GMELIN\*, IFICDB, IFIPAT,  
IFIUDB, MEDLINE, TOXCENTER, USPATFULL

(\*File contains numerically searchable property data)

$\text{H}_2\text{O}_3\text{P}-\text{NH}-\text{PO}_3\text{H}_2$

\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

77 REFERENCES IN FILE CA (1957 TO DATE)

16 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

78 REFERENCES IN FILE CAPLUS (1957 TO DATE)

3 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

REFERENCE 1: 136:402058

REFERENCE 2: 136:351957

REFERENCE 3: 135:339302

REFERENCE 4: 128:227751

REFERENCE 5: 125:268967

REFERENCE 6: 124:336819

REFERENCE 7: 123:246540

REFERENCE 8: 120:157329

REFERENCE 9: 120:98660

REFERENCE 10: 119:266619

L119 ANSWER 5 OF 15 REGISTRY COPYRIGHT 2003 ACS

RN 27216-57-5 REGISTRY

CN myo-Inositol, bis(dihydrogen phosphate) (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Inositol, bis(dihydrogen phosphate), myo- (8CI)

OTHER NAMES:

CN Inositol bisphosphate




CN Inositol diphosphate  
CN myo-Inositol bisphosphate  
CN myo-Inositol diphosphate  
FS STEREOSEARCH  
MF C6 H14 O12 P2  
CI IDS  
LC STN Files: AGRICOLA, BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CAPLUS, EMBASE,  
TOXCENTER, USPATFULL

CM 1

CRN 7664-38-2

CMF H3 O4 P




CM 2

CRN 87-89-8

CMF C6 H12 O6

Relative stereochemistry.



1117 REFERENCES IN FILE CA (1957 TO DATE)  
3 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
1117 REFERENCES IN FILE CAPLUS (1957 TO DATE)

REFERENCE 1: 138:332173  
REFERENCE 2: 138:281373  
REFERENCE 3: 138:269332  
REFERENCE 4: 137:213385  
REFERENCE 5: 136:368759  
REFERENCE 6: 136:245048  
REFERENCE 7: 135:367189  
REFERENCE 8: 135:339302  
REFERENCE 9: 134:352406

REFERENCE 10: 134:261334

L119 ANSWER 6 OF 15 REGISTRY COPYRIGHT 2003 ACS

RN 27121-72-8 REGISTRY

CN myo-Inositol, tetrakis(dihydrogen phosphate) (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Inositol tetraphosphate (6CI, 7CI)

CN Inositol, tetrakis(dihydrogen phosphate), myo- (8CI)

OTHER NAMES:

CN IP4

CN myo-Inositol tetraphosphate

FS STEREOSEARCH

MF C6 H16 O18 P4


CI IDS, COM

LC STN Files: AGRICOLA, BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CAOLD, CAPLUS, EMBASE, IFICDB, IFIPAT, IFIUDB, TOXCENTER, USPATFULL

CM 1

CRN 7664-38-2

CMF H3 O4 P




CM 2

CRN 87-89-8

CMF C6 H12 O6

Relative stereochemistry.



354 REFERENCES IN FILE CA (1957 TO DATE)

9 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

355 REFERENCES IN FILE CAPLUS (1957 TO DATE)

10 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

REFERENCE 1: 138:237164

REFERENCE 2: 138:186734

REFERENCE 3: 137:273159

REFERENCE 4: 137:213385

REFERENCE 5: 137:19668

REFERENCE 6: 136:368833

REFERENCE 7: 136:368759

REFERENCE 8: 136:366189

REFERENCE 9: 136:324404

REFERENCE 10: 136:320864

L119 ANSWER 7 OF 15 REGISTRY COPYRIGHT 2003 ACS

RN 25663-09-6 REGISTRY

CN myo-Inositol, pentakis(dihydrogen phosphate) (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Inositol pentaphosphate (7CI)

CN Inositol, pentakis(dihydrogen phosphate), myo- (8CI)


DR 10072-58-9, 53861-65-7, 56688-72-3, 90080-18-5, 27214-04-6

MF C6 H17 O21 P5

CI COM

LC STN Files: AGRICOLA, ANABSTR, BEILSTEIN\*, BIOBUSINESS, BIOSIS,  
BIOTECHNO, CA, CANCERLIT, CAOLD, CAPLUS, EMBASE, MEDLINE, TOXCENTER,  
USPATFULL

(\*File contains numerically searchable property data)



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

244 REFERENCES IN FILE CA (1957 TO DATE)

6 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

245 REFERENCES IN FILE CAPLUS (1957 TO DATE)

6 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

REFERENCE 1: 138:270607

REFERENCE 2: 138:249615

REFERENCE 3: 138:237164

REFERENCE 4: 138:13575

REFERENCE 5: 137:278170

REFERENCE 6: 137:262200

REFERENCE 7: 137:213385


REFERENCE 8: 137:19668

REFERENCE 9: 136:368833

REFERENCE 10: 136:324404

L119 ANSWER 8 OF 15 REGISTRY COPYRIGHT 2003 ACS  
 RN 25612-73-1 REGISTRY  
 CN 5'-Adenylic acid, monoanhydride with imidodiphosphoric acid (8CI, 9CI)  
 (CA INDEX NAME)  
 OTHER CA INDEX NAMES:  
 CN Imidodiphosphoric acid, monoanhydride with 5'-adenylic acid (8CI)  
 OTHER NAMES:  
 CN .beta.,.gamma.-Imido-ATP  
 CN .beta.,.gamma.-Imino-adenosine 5'-triphosphate  
 CN .beta.,.gamma.-Imino-ATP  
 CN 5'-Adenylyl (.beta.,.gamma.-imidodiphosphate)  
 CN 5'-Adenylyl imidodiphosphate  
 CN Adenosine .beta.,.gamma.-imidotriphosphate  
 CN Adenosine 5'-(.beta.,.gamma.-imidotriphosphate)  
 CN Adenosine 5'-(.beta.,.gamma.-iminotriphosphate)  
 CN Adenylyl .beta.,.gamma.-imidodiphosphate  
 CN Adenylyl imidodiphosphate  
 CN AMP-PNP  
 FS STEREOSEARCH  
 DR 27752-13-2, 33055-51-5  
 MF C10 H17 N6 O12 P3  
 CI COM  
 LC STN Files: AGRICOLA, BEILSTEIN\*, BIOBUSINESS, BIOSIS, BIOTECHNO, CA,  
 CANCERLIT, CAPLUS, DDFU, DRUGU, EMBASE, IFICDB, IFIPAT, IFIUDB, MEDLINE,  
 TOXCENTER, USPATFULL  
 (\*File contains numerically searchable property data)

Absolute stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1051 REFERENCES IN FILE CA (1957 TO DATE)  
 72 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
 1053 REFERENCES IN FILE CAPLUS (1957 TO DATE)

REFERENCE 1: 138:316772

REFERENCE 2: 138:316723

REFERENCE 3: 138:300376

REFERENCE 4: 138:299471

REFERENCE 5: 138:297992

REFERENCE 6: 138:283198

REFERENCE 7: 138:233846

REFERENCE 8: 138:200989

REFERENCE 9: 138:200988

REFERENCE 10: 138:166561

L119 ANSWER 9 OF 15 REGISTRY COPYRIGHT 2003 ACS

RN 14127-68-5 REGISTRY

CN Triphosphate (8CI, 9CI) (CA INDEX NAME)

OTHER NAMES:

CN Triphosphate (P3O105-)

CN Triphosphate(5-)

CN Tripolyphosphate

CN Tripolyphosphate (P3O105-) ion

FS 3D CONCORD

MF 010 P3


CI COM

LC STN Files: ANABSTR, BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CAPLUS, CASREACT, CEN, CHEMLIST, CIN, EMBASE, GMELIN\*, IFICDB, IFIPAT, IFIUDB, IPA, PIRA, PROMT, TOXCENTER, USPAT2, USPATFULL

(\*File contains numerically searchable property data)

Other Sources: EINECS\*\*

(\*\*Enter CHEMLIST File for up-to-date regulatory information)



585 REFERENCES IN FILE CA (1957 TO DATE)

17 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

586 REFERENCES IN FILE CAPLUS (1957 TO DATE)

REFERENCE 1: 138:344575

REFERENCE 2: 138:344420

REFERENCE 3: 138:333523

REFERENCE 4: 138:305821

REFERENCE 5: 138:291475

REFERENCE 6: 138:283141

REFERENCE 7: 138:258954

REFERENCE 8: 138:210064

REFERENCE 9: 138:175703

REFERENCE 10: 138:165498


L119 ANSWER 10 OF 15 REGISTRY COPYRIGHT 2003 ACS

RN 13598-36-2 REGISTRY

CN Phosphonic acid (6CI, 7CI, 8CI, 9CI) (CA INDEX NAME)

OTHER NAMES:

CN Dihydroxyphosphine oxide  
 CN Phosphorous acid  
 MF H3 O3 P  
 CI COM  
 LC STN Files: AGRICOLA, ANABSTR, AQUIRE, BIOBUSINESS, BIOSIS, BIOTECHNO,  
 CA, CAOLD, CAPLUS, CASREACT, CBNB, CEN, CHEMCATS, CHEMINFORMRX,  
 CHEMLIST, CIN, CSCHM, CSNB, DETHERM\*, DIPPR\*, EMBASE, IFICDB, IFIPAT,  
 IFIUDB, MRCK\*, MSDS-OHS, NIOSHTIC, PDLCOM\*, PIRA, PROMT, RTECS\*,  
 TOXCENTER, TULSA, USPAT2, USPATFULL, VTB  
 (\*File contains numerically searchable property data)  
 Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*  
 (\*\*Enter CHEMLIST File for up-to-date regulatory information)



\*\*\* FRAGMENT DIAGRAM IS INCOMPLETE \*\*\*

6229 REFERENCES IN FILE CA (1957 TO DATE)  
 3012 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
 6250 REFERENCES IN FILE CAPLUS (1957 TO DATE)  
 8 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

REFERENCE 1: 138:358492  
 REFERENCE 2: 138:358405  
 REFERENCE 3: 138:356459  
 REFERENCE 4: 138:355998  
 REFERENCE 5: 138:355997  
 REFERENCE 6: 138:355994  
 REFERENCE 7: 138:355117  
 REFERENCE 8: 138:354783  
 REFERENCE 9: 138:354136  
 REFERENCE 10: 138:352686

L119 ANSWER 11 OF 15 REGISTRY COPYRIGHT 2003 ACS


RN 7664-38-2 REGISTRY

CN Phosphoric acid (7CI, 8CI, 9CI) (CA INDEX NAME)

OTHER NAMES:

CN 3M Etching Liquid  
 CN Amberphos 54  
 CN C 134  
 CN C 134 (acid)  
 CN C 434  
 CN C 434 (acid)  
 CN Conditioner 36  
 CN Decon 4512  
 CN E 338  
 CN EVITs  
 CN HQ 54  
 CN K-etchant  
 CN Kefo  
 CN Mikro Klene DF

CN Orthophosphoric acid  
 CN Panavia Etching Agent  
 CN Sonac  
 CN SPA 2  
 CN SPA 2 (catalyst)  
 CN TG 434  
 CN Total Etch  
 CN Ultra-Etch Gel  
 CN Ultraetch  
 CN Uni-Etch  
 CN WC-Reiniger  
 FS 3D CONCORD  
 DR 28602-75-7, 178560-73-1  
 MF H3 O4 P  
 CI COM  
 LC STN Files: AGRICOLA, ANABSTR, AQUIRE, BIOBUSINESS, BIOSIS, BIOTECHNO,  
 CA, CABA, CANCERLIT, CAOLD, CAPLUS, CASREACT, CBNB, CEN, CHEMCATS,  
 CHEMINFORMRX, CHEMLIST, CIN, CSCHEM, CSNB, DDFU, DETHERM\*, DIOGENES,  
 DIPPR\*, DRUGU, EMBASE, ENCOMPLIT, ENCOMPLIT2, ENCOMPPAT, ENCOMPPAT2,  
 GMELIN\*, HSDB\*, IFICDB, IFIPAT, IFIUDB, IPA, MEDLINE, MRCK\*, MSDS-OHS,  
 NAPRALERT, NIOSHTIC, PDLCOM\*, PIRA, PROMT, RTECS\*, SPECINFO, TOXCENTER,  
 TULSA, ULIDAT, USAN, USPAT2, USPATFULL, VETU, VTB  
 (\*File contains numerically searchable property data)  
 Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*  
 (\*\*Enter CHEMLIST File for up-to-date regulatory information)



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*


52117 REFERENCES IN FILE CA (1957 TO DATE)  
 7086 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
 52174 REFERENCES IN FILE CAPLUS (1957 TO DATE)  
 1 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

REFERENCE 1: 138:361751  
 REFERENCE 2: 138:361709  
 REFERENCE 3: 138:361656  
 REFERENCE 4: 138:361639  
 REFERENCE 5: 138:361638  
 REFERENCE 6: 138:361419  
 REFERENCE 7: 138:361300  
 REFERENCE 8: 138:360392  
 REFERENCE 9: 138:360306  
 REFERENCE 10: 138:359513

RN 2809-21-4 REGISTRY  
CN Phosphonic acid, (1-hydroxyethylidene)bis- (9CI) (CA INDEX NAME)  
OTHER CA INDEX NAMES:  
CN Phosphonic acid, (1-hydroxyethylidene)di- (8CI)  
OTHER NAMES:  
CN (1-Hydroxyethylidene)-1,1-bis(phosphonic acid)  
CN (1-Hydroxyethylidene)-1,1-diphosphonic acid  
CN (1-Hydroxyethylidene)bisphosphonic acid  
CN (1-Hydroxyethylidene)diphosphonic acid  
CN 1-Hydroxy-1,1-diphosphonoethane  
CN 1-Hydroxyethane-1,1-bisphosphonic acid  
CN 1-Hydroxyethane-1,1-diphosphonic acid  
CN 1-Hydroxyethane-1,1-diylldiphosphonic acid  
CN 1-Hydroxyethanediphosphonic acid  
CN 1-Hydroxyethylidene-1,1'-diphosphonic acid  
CN 1000SL  
CN Acetodiphosphonic acid  
CN Anti Cal 5  
CN Briquest ADPA 60A  
CN Cublen K 60  
CN Defloc EH 06  
CN Dequest 16  
CN Dequest 2010  
CN Dequest 2010CS  
CN Dequest 2015  
CN Dequest Z 010  
CN EHDP  
CN Ethane-1-hydroxy-1,1-diphosphonic acid  
CN Etidronic acid  
CN Ferriox 115  
CN Ferriox CY 115  
CN Ferrofos 510  
CN HDEPA  
CN HEDP  
CN Hydroxyethanediphosphonic acid  
CN Ksidifon  
CN Lonza 106  
CN Mascol P 210  
CN Masquol P 210  
CN Mayoquest 1500  
CN OEDF  
CN OEDP  
CN Oxyethylidenediphosphonic acid  
CN RP 61  
CN Sequion 10H  
CN Sone 16  
CN Tecquest 360  
CN Terpil SL  
CN Turpinal SL  
CN Turpinal SLR  
CN Unihib 106  
CN Wayplex  
CN Xidiphone  
CN Xydiphone  
FS 3D CONCORD  
DR 129130-42-3, 51888-66-5, 66216-98-6, 138360-84-6, 85985-26-8, 86159-18-4,  
100511-44-2, 106908-76-3, 192526-55-9, 303177-33-5  
MF C2 H8 O7 P2  
CI COM  
LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, BEILSTEIN\*, BIOBUSINESS, BIOSIS,  
BIOTECHNO, CA, CABA, CANCERLIT, CAOLD, CAPLUS, CASREACT, CBNB, CEN,  
CHEMCATS, CHEMLIST, CIN, CSCHEM, DDFU, DETHERM\*, DRUGNL, DRUGU, EMBASE,  
GMELIN\*, HSDB\*, IFICDB, IFIPAT, IFIUDB, IPA, MEDLINE, MRCK\*, MSDS-OHS,



PHAR, PIRA, PROMT, RTECS\*, TOXCENTER, USAN, USPAT2, USPATFULL, VETU  
(\*File contains numerically searchable property data)  
Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*, WHO  
(\*\*Enter CHEMLIST File for up-to-date regulatory information)



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

3820 REFERENCES IN FILE CA (1957 TO DATE)  
433 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
3828 REFERENCES IN FILE CAPLUS (1957 TO DATE)  
5 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

REFERENCE 1: 138:359460  
REFERENCE 2: 138:355337  
REFERENCE 3: 138:350529  
REFERENCE 4: 138:339805  
REFERENCE 5: 138:331682  
REFERENCE 6: 138:326535  
REFERENCE 7: 138:323076  
REFERENCE 8: 138:322081  
REFERENCE 9: 138:311591  
REFERENCE 10: 138:308992

L119 ANSWER 13 OF 15 REGISTRY COPYRIGHT 2003 ACS  
RN 2466-09-3 REGISTRY

CN Diphosphoric acid (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Pyrophosphoric acid (8CI)

OTHER NAMES:

CN Diphosphoric acid (H4P2O7)


FS 3D CONCORD

DR 133883-41-7

MF H4 O7 P2

CI COM

LC STN Files: AGRICOLA, BIOBUSINESS, BIOSIS, CA, CAOLD, CAPLUS, CASREACT,  
CHEMCATS, CHEMLIST, CIN, CSCHEM, DDFU, DETHERM\*, DRUGU, EMBASE, GMELIN\*,  
HSDB\*, IFICDB, IFIPAT, IFIUDB, IPA, MEDLINE, MRCK\*, MSDS-OHS, NIOSHTIC,  
PIRA, PROMT, RTECS\*, TOXCENTER, TULSA, USPAT2, USPATFULL, VTB  
(\*File contains numerically searchable property data)  
Other Sources: EINECS\*\*  
(\*\*Enter CHEMLIST File for up-to-date regulatory information)



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

2679 REFERENCES IN FILE CA (1957 TO DATE)  
 583 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
 2683 REFERENCES IN FILE CAPLUS (1957 TO DATE)  
 20 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

REFERENCE 1: 138:356235  
 REFERENCE 2: 138:352686  
 REFERENCE 3: 138:350389  
 REFERENCE 4: 138:348010  
 REFERENCE 5: 138:340726  
 REFERENCE 6: 138:338001  
 REFERENCE 7: 138:336931  
 REFERENCE 8: 138:330038  
 REFERENCE 9: 138:311591  
 REFERENCE 10: 138:308651

L119 ANSWER 14 OF 15 REGISTRY COPYRIGHT 2003 ACS

RN 83-86-3 REGISTRY

CN myo-Inositol, hexakis(dihydrogen phosphate) (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Inositol, hexakis(dihydrogen phosphate), myo- (8CI)

OTHER NAMES:

CN Alkalover

CN D-myo-Inositol-1,2,3,4,5,6-hexaphosphate

CN Fytic acid

CN Inositol 1,2,3,4,5,6-hexakisphosphate

CN Inositol hexakis(phosphate)

CN Inositol hexaphosphate

CN IP6

CN meso-Inositol hexaphosphate

CN myo-Inositol hexakis(phosphate)

CN myo-Inositol hexaphosphate

CN Phytic acid

FS STEREOSEARCH

DR 50762-79-3, 78039-41-5

MF C6 H18 O24 P6


CI COM

LC

STN Files: AGRICOLA, ANABSTR, BEILSTEIN\*, BIOBUSINESS, BIOSIS,  
 BIOTECHNO, CA, CABA, CANCERLIT, CAOLD, CAPLUS, CASREACT, CBNB, CEN,  
 CHEMCATS, CHEMLIST, CIN, CSCHEM, DDFU, DRUGU, EMBASE, GMELIN\*, IFICDB,  
 IFIPAT, IFIUDB, IPA, MEDLINE, MRCK\*, MSDS-OHS, NAPRALERT, NIOSHTIC,  
 PHAR, PIRA, PROMT, RTECS\*, TOXCENTER, TULSA, USAN, USPAT2, USPATFULL,  
 VETU

(\*File contains numerically searchable property data)  
Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*, WHO  
(\*\*Enter CHEMLIST File for up-to-date regulatory information)

Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

4983 REFERENCES IN FILE CA (1957 TO DATE)  
231 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
4996 REFERENCES IN FILE CAPLUS (1957 TO DATE)  
5 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

REFERENCE 1: 138:356000  
REFERENCE 2: 138:353034  
REFERENCE 3: 138:339568  
REFERENCE 4: 138:336853  
REFERENCE 5: 138:336844  
REFERENCE 6: 138:336746  
REFERENCE 7: 138:336707  
REFERENCE 8: 138:336577  
REFERENCE 9: 138:334429  
REFERENCE 10: 138:329033

L119 ANSWER 15 OF 15 REGISTRY COPYRIGHT 2003 ACS

RN 52-68-6 REGISTRY

CN Phosphonic acid, (2,2,2-trichloro-1-hydroxyethyl)-, dimethyl ester (6CI, 8CI, 9CI) (CA INDEX NAME)

OTHER NAMES:

CN (.+-.)-Trichlorfon  
CN 1-Hydroxy-2,2,2-trichloroethylphosphonate-O,O-dimethyl ester  
CN Aerol 1 (pesticide)  
CN Agroforotox  
CN Anthon  
CN BAY-a 9826  
CN BAY-L 1359  
CN Bayer L 13/59  
CN Bayer L 1359  
CN Chlorak  
CN Chlorofos  
CN Chloroftalm  
CN Chlorophos  
CN Chlorophthalm  
CN Chloroxyphos

CN Combot  
 CN Danex  
 CN DEP  
 CN DEP (pesticide)  
 CN Depthon  
 CN DETF  
 CN Dimethyl (2,2,2-trichloro-1-hydroxyethyl)phosphonate  
 CN Dimethyl 1-hydroxy-2,2,2-trichloroethylphosphonate  
 CN Dimetox  
 CN Dioxaphos  
 CN Dipterex  
 CN Dipterex 50  
 CN Dipterex 500  
 CN Dipterex SL  
 CN Dipterex WP 80  
 CN Diptevur  
 CN Ditrifon  
 CN Ditrifon 50  
 CN Dylox  
 CN Dyrex  
 CN Dyvon  
 CN ENT 19,763  
 CN Flibol E  
 CN Fliegenteller  
 CN Forotox  
 CN Foschlor  
 CN Foschlor R  
 CN Foschlor R 50  
 CN Hypodermacid  
 CN Loisol  
 CN Masoten  
 CN Mazoten  
 CN Methyl chlorophos  
 CN Metrifonate  
 CN Metriphosphate

ADDITIONAL NAMES NOT AVAILABLE IN THIS FORMAT - Use FCN, FIDE, or ALL for  
 DISPLAY


FS 3D CONCORD  
 DR 56042-25-2, 66758-31-4, 50924-44-2, 37333-09-8  
 MF C4 H8 Cl3 O4 P  
 CI COM

LC STN Files: ADISINSIGHT, ADISNEWS, AGRICOLA, ANABSTR, AQUIRE, BEILSTEIN\*,  
 BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CABA, CANCERLIT, CAOLD, CAPLUS,  
 CASREACT, CBNB, CHEMCATS, CHEMLIST, CIN, CSCHEM, CSNB, DDFU, DETHERM\*,  
 DIOGENES, DRUGNL, DRUGPAT, DRUGU, DRUGUPDATES, EMBASE, HSDB\*, IFICDB,  
 IFIPAT, IFIUDB, IPA, MEDLINE, MRCK\*, MSDS-OHS, NIOSHTIC, PHAR,  
 PHARMASEARCH, PIRA, PROMT, RTECS\*, SPECINFO, SYNTHLINE, TOXCENTER,  
 ULIDAT, USAN, USPAT2, USPATFULL, VETU, VTB

(\*File contains numerically searchable property data)

Other Sources: EINECS\*\*, WHO

(\*\*Enter CHEMLIST File for up-to-date regulatory information)



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

5169 REFERENCES IN FILE CA (1957 TO DATE)  
20 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
5171 REFERENCES IN FILE CAPLUS (1957 TO DATE)  
297 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

REFERENCE 1: 138:333151  
REFERENCE 2: 138:299194  
REFERENCE 3: 138:297496  
REFERENCE 4: 138:287669  
REFERENCE 5: 138:286302  
REFERENCE 6: 138:282719  
REFERENCE 7: 138:281016  
REFERENCE 8: 138:267208  
REFERENCE 9: 138:267201  
REFERENCE 10: 138:251130

=> fil hcaplus

FILE 'HCAPLUS' ENTERED AT 09:26:19 ON 04 JUN 2003  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 4 Jun 2003 VOL 138 ISS 23  
FILE LAST UPDATED: 3 Jun 2003 (20030603/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d all tot 1117

L117 ANSWER 1 OF 10 HCAPLUS COPYRIGHT 2003 ACS  
AN 2002:682100 HCAPLUS  
TI Inactivation of the human brain muscarinic acetylcholine receptor by oxidative damage catalyzed by a low molecular weight endogenous inhibitor from Alzheimer's brain is prevented by **pyrophosphate** analogs, bioflavonoids and other antioxidants  
AU **Fawcett, John R.**; Bordayo, Elizabeth Z.; Jackson, Kathy; Liu, Howard; Peterson, Jennifer; Svitak, Aleta; **Frey, William H., II**  
CS The Alzheimer's Research Center, Regions Hospital, **HealthPartners** Research Foundation, St. Paul, MN, 55101-2595, USA  
SO Brain Research (2002), 950(1,2), 10-20

CODEN: BRREAP; ISSN: 0006-8993

PB Elsevier Science B.V.

DT Journal

LA English

CC 1-11 (Pharmacology)

- AB Oxidative stress has been implicated as a contributing factor to neurodegeneration in Alzheimer's disease. An endogenous, low mol. wt. (LMW) inhibitor from Alzheimer's brain inactivates the human brain muscarinic acetylcholine receptor (mAChR). The inhibitor prevents agonist and antagonist binding to the mAChR as assessed by radioligand binding studies. The LMW endogenous inhibitor, which has components with mol. wts. between 100 and 1000 Da, requires dissolved oxygen and glutathione. Prevention of inactivation of the mAChR with peroxidase suggests that the LMW endogenous inhibitor generates peroxide. Heme, previously shown to be present in the LMW endogenous inhibitor, also inactivates the mAChR in the presence of peroxide. Free radical damage to the muscarinic receptor by the endogenous inhibitor can be prevented through the use of naturally occurring antioxidants including bilirubin, biliverdin, carnosol, myricetin and quercetin. In addn., **pyrophosphate**, **imidodiphosphate**, **bisphosphonates** and related compds. also protect the muscarinic receptor from free radical damage. Inactivation of the mAChR by the LMW endogenous inhibitor is likely to be a factor in the continual decline of Alzheimer's patients, even those taking acetylcholinesterase inhibitors. Natural antioxidants and **pyrophosphate** analogs may improve the effectiveness of acetylcholinesterase inhibitors and prove useful in the treatment and prevention of Alzheimer's disease since the muscarinic acetylcholine receptor is required for memory, and decreased cholinergic function is a crit. deficit in Alzheimer's disease.
- ST muscarinic acetylcholine receptor oxidative damage Alzheimer disease **pyrophosphate** analog; neuroprotectant bioflavonoid antioxidant  
Alzheimer disease **pyrophosphate** analog muscarinic receptor
- IT Flavonoids  
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(bioflavonoids; inactivation of human brain muscarinic acetylcholine receptor by oxidative damage catalyzed by a low mol. wt. endogenous inhibitor from Alzheimer's brain is prevented by **pyrophosphate** analogs, bioflavonoids and other antioxidants)
- IT **Anti-Alzheimer's agents**  
Brain  
Cognition enhancers  
Human  
Memory, biological  
Oxidative stress, biological  
(inactivation of human brain muscarinic acetylcholine receptor by oxidative damage catalyzed by a low mol. wt. endogenous inhibitor from Alzheimer's brain is prevented by **pyrophosphate** analogs, bioflavonoids and other antioxidants)
- IT Muscarinic receptors  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(inactivation of human brain muscarinic acetylcholine receptor by oxidative damage catalyzed by a low mol. wt. endogenous inhibitor from Alzheimer's brain is prevented by **pyrophosphate** analogs, bioflavonoids and other antioxidants)
- IT Antioxidants  
(natural; inactivation of human brain muscarinic acetylcholine receptor by oxidative damage catalyzed by a low mol. wt. endogenous inhibitor from Alzheimer's brain is prevented by **pyrophosphate** analogs, bioflavonoids and other antioxidants)
- IT 114-25-0, Biliverdin 117-39-5, Quercetin 529-44-2, Myricetin 635-65-4, Bilirubin 2466-09-3, Diphosphoric acid 5957-80-2, Carnosol 14127-68-5, Tripolyphosphate

25612-73-1 27590-04-1, Imidodiphosphoric acid

34273-04-6 40391-99-9 89771-93-7, Bilirubin ditaurate

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL  
(Biological study); USES (Uses)

(inactivation of human brain muscarinic acetylcholine receptor by  
oxidative damage catalyzed by a low mol. wt. endogenous inhibitor from  
Alzheimer's brain is prevented by **pyrophosphate** analogs,  
bioflavonoids and other antioxidants)

RE.CNT 47 THERE ARE 47 CITED REFERENCES AVAILABLE FOR THIS RECORD  
RE

- (1) Albrow, P; J Inorg Biochem 1986, V27, P191 HCAPLUS
- (2) American Psychiatric Association; Diagnostic and Statistical Manual of  
Mental Disorders 4th Edition 2000
- (3) Boyer, J; Biochem Biophys Res Commun 1986, V134, P172 HCAPLUS
- (4) Broody, K; Br J Radiol 1976, V49, P267
- (5) Budavari, S; An Encyclopedia of Chemicals, Drugs, and Biologicals 12th  
Edition 1996, P205
- (6) Buxbaum, J; Proc Natl Acad Sci 1992, V89, P10075 HCAPLUS
- (7) Commenges, D; Eur J Epidemiol 2000, V16, P357 HCAPLUS
- (8) Coyle, J; Biol Psychiatry 2001, V49, P289 HCAPLUS
- (9) Davies, P; Lancet 1976, V2, P1403 MEDLINE
- (10) De Souza, E; Advances in Clinical and Basic Research 1993, P539
- (11) Farlow, M; Eur Neurol 2000, V44, P236 HCAPLUS
- (12) Fields, J; J Biol Chem 1978, V253, P3251 HCAPLUS
- (13) Frey, W; Brain Res 1994, V655, P153 HCAPLUS
- (14) Frey, W; Brain Res 1996, V714, P87 HCAPLUS
- (15) Genis, I; J Neurochem 1999, V72, P206 HCAPLUS
- (16) Hake, A; Cleve Clin J Med 2001, V68, P608 MEDLINE
- (17) Hansen, T; Pediatric Res 2001, V49, P203 HCAPLUS
- (18) Hortobagyi, G; J Clin Oncol 1998, V16, P2038 MEDLINE
- (19) Kanowski, S; Pharmacopsychiatry 1996, V29, P47 MEDLINE
- (20) Katzman, R; Basic Neurochemistry 1989, P827
- (21) Khachaturian, Z; Arch Neurol 1985, V42, P1097 MEDLINE
- (22) Lander, C; Exp Neurol 1999, V158, P451
- (23) Marks, M; Mol Pharmacol 1982, V22, P554 HCAPLUS
- (24) Mirra, S; Neurology 1991, V41, P479 MEDLINE
- (25) Mohs, R; Neurology 2001, V57, P481 HCAPLUS
- (26) Monji, A; J Neurochem 2001, V77, P1425 HCAPLUS
- (27) Nitsch, R; Ann NY Acad Sci P759
- (28) Nitsch, R; Ann NY Acad Sci 1996, V777, P175 HCAPLUS
- (29) Nunomura, A; J Neuropathol Exp Neurol 2001, V60 HCAPLUS
- (30) Ocken, B; Arch Neurol 1998, V55, P1409
- (31) Pavia, J; Fundam Clin Pharmacol 1998, V12, P473 HCAPLUS
- (32) Premkumar, D; J Neurochem 1995, V65, P1399 HCAPLUS
- (33) Rhine, W; J Perinatol 2001, V19, P206
- (34) Rodriguez-Puertas, R; Synapse 1997, V26, P341 HCAPLUS
- (35) Rottkamp, C; Alzheimer Dis Assoc Dis 2000, V14, P562 HCAPLUS
- (36) Sadot, E; J Neurochem 1996, V66, P877 HCAPLUS
- (37) Sadrzadeh, S; J Biol Chem 1984, V259, P14354
- (38) Schipper, H; Ann Neurol 1995, V37, P758 MEDLINE
- (39) Smith, M; Am J Pathol 1994, V145, P42 MEDLINE
- (40) Smith, M; Anti Redox Signalling 2000, V2, P413 HCAPLUS
- (41) Smith, P; Ann Biochem 1985, V150, P76 HCAPLUS
- (42) Soto-Otero, R; Life Sci 2001, V69, P879 HCAPLUS
- (43) Subramanian, G; JNM/Radiochem Radiopharmaceutics 1975, V16, P1137 HCAPLUS
- (44) Takahashi, M; Neuron 2000, V28, P461 HCAPLUS
- (45) Tsujimoto, Y; Gen Pharmacol 1998, V31, P405 HCAPLUS
- (46) Venters, H; Brain Res 1997, V764, P93
- (47) Whitehouse, P; Science 1982, V215, P1237 MEDLINE

L117 ANSWER 2 OF 10 HCAPLUS COPYRIGHT 2003 ACS

AN 2001:833023 HCAPLUS

DN 135:376738

TI Compounds and methods for modulating cerebral **amyloid** angiopathy  
using inhibitors of an **amyloid** .beta. peptide  
IN Green, Allan M.; Gervais, Francine  
PA Neurochem, Inc., Can.  
SO PCT Int. Appl., 68 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K

CC 63-6 (Pharmaceuticals)

Section cross-reference(s): 1

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001085093	A2	20011115	WO 2000-IB2078	20001222 <--
	WO 2001085093	A3	20020829		
	WO 2001085093	C2	20020926		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
	AU 2001084313	A5	20011120	AU 2001-84313	20001222 <--
	EP 1251837	A2	20021030	EP 2000-993855	20001222 <--
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
	BR 2000016652	A	20021119	BR 2000-16652	20001222 <--
	US 2003003141	A1	20030102	US 2000-747408	20001222 <--
PRAI	US 1999-171877P	P	19991223 <--		
	WO 2000-IB2078	W	20001222		

OS MARPAT 135:376738

AB The invention provides methods of inhibiting cerebral **amyloid** angiopathy (CAA) and treating a disease state characterized by cerebral **amyloid** angiopathy, e.g., **Alzheimer's** disease, in a subject using an inhibitor of the 39-40 amino acid **amyloid** .beta. peptide (A.beta.40). The A.beta.40 inhibitor is selected from, e.g., sulfonic acid derivs., such as ethanesulfonic acid, 1,2-ethanedisulfonic acid, 1-propanesulfonic acid, 1,3-propanedisulfonic acid, 1,4-butanedisulfonic acid, 1,5-pentanedisulfonic acid, 2-aminoethanesulfonic acid, 4-hydroxy-1-butanedisulfonic acid, 1-butanedisulfonic acid, 1-decanedisulfonic acid, 2-propanedisulfonic acid, 3-pentanesulfonic acid, 4-heptanesulfonic acid, etc., and pharmaceutically acceptable salts thereof or from from phosphonic acid derivs., such as diethylphosphonoacetic acid, phenylphosphonic acid, 3-aminopropylphosphonic acid, propylphosphonic acid, etc. The compds. are formulated in a dispersion system, a liposome formulation, or microspheres using a polymeric matrix. The polymeric matrix is selected from natural polymers, such as albumin, alginate, cellulose derivs., collagen, fibrin, gelatin, and polysaccharides, or synthetic polymers such as polyesters, polyethylene glycol, poloxamers, and polyanhydrides. For example, the ability of compds. of the invention to inhibit CAA was measured in 9 wk old hAPP transgenic mice treated with two different concns. of a compd. of the present invention, 3-amino-1-propanedisulfonic acid sodium salt, 100 and 30 mg/kg. Mice were administered the compd. for 8 wk, after which they were sacrificed and their brains were perfused and processed for histol. staining with Thioflavin S. This method may also be used as a screening method for detg. activity of a candidate compd. for inhibiting CAA. The extent of CAA in brain sections obtained from these animals was qual. detd. following staining. The results indicate that the test compd. was



- effective in (i) reducing the no. of mice showing CAA, and (ii) showing an effect on the severity of the deposition seen in the brain vasculature of these animals.
- ST **amyloid** peptide inhibitor cerebral **amyloid** angiopathy;  
sulfonate phosphonate **amyloid** peptide inhibitor
- IT Brain, disease  
(**amyloid** angiopathy; inhibitors of **amyloid** .beta.  
peptide for modulating cerebral **amyloid** angiopathy)
- IT Blood vessel  
(endothelium; inhibitors of **amyloid** .beta. peptide for  
modulating cerebral **amyloid** angiopathy)
- IT Brain, disease  
(hemorrhagic stroke; inhibitors of **amyloid** .beta. peptide for  
modulating cerebral **amyloid** angiopathy)
- IT **Amyloidosis**  
(hereditary, cerebral hemorrhage type, Dutch type; inhibitors of  
**amyloid** .beta. peptide for modulating cerebral **amyloid**  
angiopathy)
- IT Polyesters, biological studies  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(hydroxycarboxylic acid-based, matrixes; inhibitors of **amyloid**  
.beta. peptide for modulating cerebral **amyloid** angiopathy)
- IT **Anti-Alzheimer's agents**  
Blood vessel  
Diagnosis  
Imaging agents  
Peptidomimetics  
(inhibitors of **amyloid** .beta. peptide for modulating cerebral  
**amyloid** angiopathy)
- IT Peptides, biological studies  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological  
study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES  
(Uses)  
(inhibitors of **amyloid** .beta. peptide for modulating cerebral  
**amyloid** angiopathy)
- IT Lipids, biological studies  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(inhibitors of **amyloid** .beta. peptide for modulating cerebral  
**amyloid** angiopathy)
- IT Polyesters, biological studies  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(lactic acid-based, matrixes; inhibitors of **amyloid** .beta.  
peptide for modulating cerebral **amyloid** angiopathy)
- IT Drug delivery systems  
(liposomes, multivesicular; inhibitors of **amyloid** .beta.  
peptide for modulating cerebral **amyloid** angiopathy)
- IT Drug delivery systems  
(liqs., dispersions; inhibitors of **amyloid** .beta. peptide for  
modulating cerebral **amyloid** angiopathy)
- IT Albumins, biological studies  
Biopolymers  
Collagens, biological studies  
Fibrins  
Gelatins, biological studies  
Polyanhydrides  
Polymers, biological studies  
Polyoxyalkylenes, biological studies  
Polysaccharides, biological studies  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(matrixes; inhibitors of **amyloid** .beta. peptide for  
modulating cerebral **amyloid** angiopathy)
- IT Drug delivery systems  
(microspheres; inhibitors of **amyloid** .beta. peptide for

modulating cerebral **amyloid** angiopathy)

IT Capillary vessel  
(pericyte; inhibitors of **amyloid** .beta. peptide for  
modulating cerebral **amyloid** angiopathy)

IT Cell death  
(prevention of; inhibitors of **amyloid** .beta. peptide for  
modulating cerebral **amyloid** angiopathy)

IT Muscle  
(smooth, blood vessel; inhibitors of **amyloid** .beta. peptide  
for modulating cerebral **amyloid** angiopathy)

IT Drug delivery systems  
(sustained-release; inhibitors of **amyloid** .beta. peptide for  
modulating cerebral **amyloid** angiopathy)

IT **Amyloid**  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(.beta.-; inhibitors of **amyloid** .beta. peptide for modulating  
cerebral **amyloid** angiopathy)

IT 81-08-3 107-35-7, 2-Aminoethanesulfonic acid 110-04-3,  
1,2-Ethanedisulfonic acid 116-63-2 149-45-1 288-94-8, 1H-Tetrazole  
594-45-6, Ethanesulfonic acid 831-59-4 860-22-0 926-39-6 993-13-5,  
Methylphosphonic acid 1068-21-9, Diethyl phosphoramidate 1071-83-6,  
N-Phosphonomethylglycine 1120-71-4 1132-61-2, 4-  
Morpholinepropanesulfonic acid 1135-40-6 1571-33-1, Phenylphosphonic  
acid 1633-83-6 2386-47-2, 1-Butanesulfonic acid 2386-54-1  
3095-95-2, Diethylphosphonoacetic acid 3687-18-1, 3-Amino-1-  
propanesulfonic acid 4408-78-0, Phosphonoacetic acid 4426-50-0  
4672-38-2, Propylphosphonic acid 4923-84-6 5117-07-7 5284-66-2,  
1-Propanesulfonic acid 5399-58-6 5652-28-8 5994-73-0 6779-09-5,  
Ethylphosphonic acid 7365-45-9 13138-33-5, 3-Aminopropylphosphonic  
acid 13419-61-9 13991-98-5 14047-23-5, (1-Aminopropyl)phosphonic  
acid 14159-48-9, 2-Propanesulfonic acid 14650-46-5 15471-17-7  
15763-57-2 18039-42-4 20283-21-0, 1-Decanesulfonic acid 21668-77-9,  
1,3-Propanedisulfonic acid 23052-80-4 23052-81-5 25331-57-1  
25595-59-9 26978-64-3, 4-Hydroxy-1-butanedisulfonic acid 27665-39-0,  
1,4-Butanedisulfonic acid 27797-35-9 31465-25-5 34159-44-9  
36585-99-6 37810-68-7 38911-09-0 **40391-99-9** 40465-65-4,  
N-Phosphonomethylglycine trisodium salt 51224-03-4 51224-04-5  
51650-30-7, 3-Pentanesulfonic acid 51762-95-9 53329-36-5 57605-13-7  
58849-79-9 60142-96-3 63585-09-1, Phosphonoformic acid trisodium salt  
71119-22-7 72217-85-7 73858-58-9 75277-39-3 76326-31-3,  
2-Amino-5-phosphonopentanoic acid 78739-01-2, D-(-)-2-Amino-4-  
phosphonobutanoic acid 79055-67-7 79055-68-8 81338-23-0  
81338-24-1, L-(+)-2-Amino-7-phosphonoheptanoic acid 82283-67-8  
82283-68-9 82977-27-3 87625-44-3 88246-85-9 91357-22-1  
91586-81-1 99107-93-4 101020-77-3, 1,5-Pentanedisulfonic acid  
102805-84-5 108084-41-9 112980-83-3 117414-74-1 126253-57-4  
126453-07-4 128241-72-5 129318-43-0 131177-53-2 138199-51-6  
143018-67-1 145544-51-0 157381-42-5 168977-94-4,  
3-Aminopropyl(methyl)phosphinic acid hydrochloride 183278-21-9,  
4-Heptanesulfonic acid 183278-22-0 183278-30-0 183278-32-2  
183278-33-3 183278-34-4 183278-35-5 183278-36-6 183505-70-6  
186295-21-6 186295-22-7 188642-79-7 205248-54-0 216367-45-2  
216367-47-4 216367-53-2 216367-60-1 216367-67-8 216367-68-9  
216367-72-5 216367-73-6 216367-74-7 216367-77-0 216367-78-1  
216367-85-0 216367-87-2 216367-89-4 216367-92-9 216367-93-0  
216367-94-1 216367-99-6 216368-01-3 216368-05-7 309751-75-5  
309751-76-6 309751-77-7 331809-98-4 373644-10-1 373644-12-3  
373644-13-4 373644-14-5 373644-15-6 373644-16-7 373644-17-8  
373644-18-9 373644-19-0 373644-20-3 373644-21-4 373644-22-5  
373644-23-6 373644-24-7 373644-25-8 373644-26-9 373644-27-0  
373644-28-1 373644-29-2 373644-30-5 373644-31-6 373644-32-7  
373644-33-8 373644-34-9 373644-35-0 373644-36-1 373644-37-2  
373644-38-3 373644-39-4 373644-40-7 373644-41-8 373644-42-9

373644-43-0	373644-44-1	373644-45-2	373644-46-3	373644-47-4
373644-48-5	373644-49-6	373644-50-9	373644-51-0	373644-52-1
373644-53-2	373644-56-5	373644-58-7	373644-60-1	373644-62-3
373644-63-4	373644-64-5	373644-65-6	373644-66-7	373644-67-8
373644-68-9	373644-69-0	373644-70-3	373644-71-4	373644-72-5
373644-73-6	373644-74-7	373644-75-8	373644-76-9	373644-77-0
373644-78-1	373644-79-2	373644-80-5	373644-81-6	373644-82-7
373644-83-8	373644-84-9	373644-85-0	373644-86-1	373644-87-2
373644-88-3	373644-89-4	373644-90-7	373644-91-8	373644-92-9
373644-93-0	373644-94-1	373644-95-2	373644-96-3	373644-97-4
373644-98-5	373644-99-6	373645-00-2	373645-01-3	373645-02-4
373645-03-5	373645-04-6	373645-05-7	373645-06-8	373645-07-9
373645-08-0	373645-09-1	373645-10-4	373645-11-5	373645-12-6
373645-13-7	373645-14-8			

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(inhibitors of amyloid .beta. peptide for modulating cerebral amyloid angiopathy)

IT	373645-15-9	373645-16-0	373645-17-1	373645-18-2	373645-19-3
	373645-20-6	373645-21-7	373645-22-8	373645-23-9	373645-24-0
	373645-25-1	373645-26-2	373645-27-3	373645-28-4	373645-29-5
	373645-30-8	373645-31-9	373645-32-0	373645-33-1	373645-34-2
	373645-35-3	373645-36-4	373645-37-5	373645-38-6	373645-39-7
	373645-40-0	373645-41-1	373645-42-2	373645-43-3	373645-44-4
	373645-45-5	373645-46-6	373645-47-7	373645-48-8	373645-49-9
	373645-50-2	373645-51-3	373645-52-4	373645-53-5	373645-54-6
	373645-56-8	373645-57-9	373645-58-0	373645-59-1	373645-60-4
	374078-81-6	374078-82-7			

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(inhibitors of amyloid .beta. peptide for modulating cerebral amyloid angiopathy)

IT 9004-34-6D, Cellulose, ethers, biological studies 9005-32-7, Alginic acid 25322-68-3, Polyethylene glycol 26023-30-3, Poly[oxy(1-methyl-2-oxo-1,2-ethanediyl)] 26100-51-6, Poly(lactic acid) 34346-01-5, Glycolic acid-lactic acid copolymer 106392-12-5, Poloxamer

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(matrixes; inhibitors of amyloid .beta. peptide for modulating cerebral amyloid angiopathy)

L117 ANSWER 3 OF 10 HCAPLUS COPYRIGHT 2003 ACS

AN 2001:816459 HCAPLUS

DN 135:339302

TI Methods and compositions for enhancing cellular function through protection of tissue components

IN Frey, William H., II; Fawcett, John Randall; Thorne, Robert Gary; Chen, Xueqing

PA Healthpartners Research Foundation, USA

SO PCT Int. Appl., 77 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K031-661

ICS A61K031-6615; A61K031-662; A61K031-7084; A61K031-706; A61K031-7076

CC 1-12 (Pharmacology)

Section cross-reference(s): 2, 4, 63

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001082932	A2	20011108	WO 2001-US13931	20010430 <--
	WO 2001082932	A3	20020718		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

US 2002028786 A1 20020307 US 2001-844450 20010427 <--

EP 1278525 A2 20030129 EP 2001-930957 20010430 <--

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR

PRAI US 2000-200843P P 20000501 <--

US 2000-230263P P 20000906 <--

US 2000-233025P P 20000915 <--

US 2000-233263P P 20000918 <--

WO 2001-US13931 W 20010430 <--

OS MARPAT 135:339302

AB Methods and compns. for enhancing cellular function through protection of tissue components, such as receptors, proteins, lipids, nucleic acids, carbohydrates, hormones, vitamins, and cofactors, by administering **pyrophosphate** analogs or related compds. Preferably, the invention provides a method for protecting a muscarinic acetylcholine receptor (mAChR) an/or increasing the efficacy of and agent the directly or indirectly affects a mAChR in a subject in need thereof.

ST tissue component protection **pyrophosphate** analog; cellular function tissue protection **pyrophosphate** analog; muscarinic receptor protection **pyrophosphate** analog

IT Bone, disease

(Paget's, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Mental disorder

(affective, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Diagnosis

(agents, formulation with **pyrophosphate** derivs. of; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Nervous system

(amyotrophic lateral sclerosis, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Receptors

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(aspartate, protection of; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Infection

(bacterial, treatment of; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Gene, animal

RL: BAC (Biological activity or effector, except adverse); BSU (Biological

study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(biliverdin reductase-encoding; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Flavonoids

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(bioflavonoids; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Neurotrophic factors

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(brain-derived; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Antitumor agents

(brain; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Cytoprotective agents

(cardioprotective; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Gene, animal

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(catalase-encoding; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Nervous system

(central, disease, from aging, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Nervous system

(central, infection, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Nervous system

(cerebellar ataxia, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Lung, disease

(chronic obstructive, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Nucleosides, biological studies

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES

(Uses)

(di, 5.5'-**pyrophosphate** derivs.; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Mental disorder

(diffuse Lewy body disease, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Esophagus

(disease, achalasia, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Spinal cord

(disease, stroke, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Blood

Endocrine system

Nervous system

(disease, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Intestine, disease

(diverticulitis, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Heart, disease

(failure, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Poisoning, biological

(from metals, protection from; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Neurotrophic factors

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(glial-derived; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Gene, animal

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(heme oxygenase 1- and 2-encoding; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT Proteins, specific or class

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(heme-binding; methods and compns. for enhancing cellular function

through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

- IT Hemochromatosis  
(hereditary hemochromatosis, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Bladder  
(incontinence, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Heart, disease  
(infarction, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Animal  
Mammal (Mammalia)  
Plant (Embryophyta)  
(infection in, treatment of; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Algae  
(infection with, treatment of; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Brain, neoplasm  
(inhibitors; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Intestine, disease  
(irritable bowel syndrome, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Heart, disease  
(ischemia, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Carcinogens  
(metals, protection from poisoning from; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT **Anti-Alzheimer's agents**  
Antiarrhythmics  
Antidiabetic agents  
Antihypertensives  
Antiparkinsonian agents  
Antipsychotics  
Antitumor agents  
Anxiolytics  
Cytoprotective agents  
Drug interactions  
Gene therapy  
Genetic vectors  
Muscarinic agonists

- Muscarinic antagonists
- Nervous system agents
  - (methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Ciliary neurotrophic factor
  - Gangliosides
  - Hemopexins
  - Lipoproteins
  - Neurokinins
  - Neurotrophic factors
    - Phosphatidylserines**
  - Platelet-derived growth factors
    - Polyphosphates**
  - Thyroid hormones
  - Transforming growth factors
- RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
  - (methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Spinal cord
  - (neoplasm, inhibitors; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Plasmid vectors
  - (nerve growth factor-encoding; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Gene, animal
  - mRNA
  - RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
    - (nerve growth factor-encoding; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Nerve, disease
  - (neuropathy, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Cytoprotective agents
  - (neuroprotectants; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Proteins, specific or class
  - RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
    - (nexins, glial-derived; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Gene, animal
  - RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)



(peroxidase-encoding; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

## IT Cations

(protection from poisoning from; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

## IT Metals, biological studies

RL: ADV (Adverse effect, including toxicity); BIOL (Biological study)

(protection from poisoning from; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

## IT Oxidative stress, biological

(protection from; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

## IT 5-HT receptors

Benzodiazepine receptors

Cannabinoid receptors

Carbohydrates, biological studies

Catecholamine receptors

GABA receptors

Glutamate receptors

Glycine receptors

Growth factor receptors

Histamine receptors

Hormones, animal, biological studies

Lipids, biological studies

Muscarinic receptors

Neuropeptide receptors

Neurotransmitter receptors

Neurotrophic factor receptors

Nicotinic receptors

Nucleic acids

Opioid receptors

Proteins, general, biological studies

Purinoceptors

Receptors

Steroid receptors

Vitamins

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL

(Biological study); PROC (Process)

(protection of; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

## IT Paralysis

(pseudobulbar, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

## IT Drug delivery systems

(**pyrophosphate** analog-contg.; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

## IT Odor and Odorous substances

(receptors for, protection of; methods and compns. for enhancing cellular function through protection of tissue components such as

- muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Ion channel  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(receptors for, protection of; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Muscle, disease  
(smooth, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Antitumor agents  
(spinal cord; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Brain, disease  
Prion diseases  
(stroke, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Antisense oligonucleotides  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(to nerve growth factors; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Aging, animal  
(treatment of diseases in; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Mycosis  
(treatment of; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents).
- IT Blood vessel, disease  
Bone, disease  
Heart, disease  
Mental disorder  
Multiple sclerosis  
Schizophrenia  
Sjogren's syndrome  
(treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT Mouth  
(xerostomia, treatment; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)
- IT 9059-22-7, Heme oxygenase  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(1 and 2; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT 13598-36-2D, **Phosphonic acid**, alkylidenebis- derivs.

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); **THU (Therapeutic use)**; BIOL (Biological study); USES (Uses)

(**Bisphosphonate**; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT 9001-08-5, Cholinesterase

RL: BSU (Biological study, unclassified); BIOL (Biological study) (inhibitors; methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT 52-68-6, Metrifonate 83-86-3, **Inositol**

**hexaphosphate** 114-25-0, Biliverdin 117-39-5, Quercetin 120-73-0D, Purine, **cyclopyrophosphate** analogs 141-43-5, Ethanolamine, biological studies 153-76-4, Gallamine 289-95-2D, Pyrimidine, acyclonucleoside analogs 357-70-0, Galanthamine 529-44-2, Myricetin 635-65-4, Bilirubin, biological studies 2466-09-3,

**Diphosphoric acid** 2466-09-3D, **Diphosphoric**

**acid**, analogs 2809-21-4, **Etidronic acid**

5957-80-2, Carnosol 6893-02-3, T3 9001-05-2, Catalase 9003-99-0, Peroxidase 9004-10-8, Insulin, biological studies 9061-61-4, Nerve growth factor 9074-10-6, Biliverdin reductase 14127-68-5,

**Tripolyphosphate** 25612-73-1 25663-09-6,

**Inositol pentaphosphate** 27121-72-8,

**Inositol tetraphosphate** 27216-57-5,

**Inositol diphosphate** 27590-04-1,

**Imidodiphosphoric acid** 34273-04-6 37758-47-7, GM-1

ganglioside 40391-99-9, **Pamidronic acid**

61912-98-9, Insulin-like growth factor 62031-54-3, Fibroblast growth factor 62229-50-9, Epidermal growth factor 67763-96-6, Insulin-like growth factor 1 67763-97-7, Insulin-like growth factor 2

**85166-31-0, Inositol triphosphate**

106096-92-8, Acidic fibroblast growth factor 106096-93-9, Basic

fibroblast growth factor 120014-06-4, Donepezil 123441-03-2,

Rivastigmine 130939-66-1, Neurotrophin 3 131986-45-3, Xanomeline

140698-57-3, Activity-dependent neurotrophic factor 143375-33-1,

Neurotrophin 4 148499-03-0, Neurotrophin 5

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); **THU (Therapeutic use)**; BIOL (Biological study); USES (Uses)

(methods and compns. for enhancing cellular function through protection of tissue components such as muscarinic receptors by administering **pyrophosphate** analogs and combination with other agents)

IT 7439-89-6, Iron, biological studies 7439-92-1, Lead, biological studies 7439-97-6, Mercury, biological studies 7440-02-0, Nickel, biological studies 7440-38-2, Arsenic, biological studies 7440-43-9, Cadmium, biological studies 7440-47-3, Chromium, biological studies 7440-48-4, Cobalt, biological studies 7440-50-8, Copper, biological studies 7440-62-2, Vanadium, biological studies 14280-50-3, Lead, ion (Pb+2), biological studies 14302-87-5, Mercuric ion, biological studies 15158-11-9, Cupric ion, biological studies 15438-31-0, Ferrous ion, biological studies 22537-48-0, Cadmium, ion (Cd+2), biological studies 22541-54-4, Arsenic, ion (As+3), biological studies

RL: ADV (Adverse effect, including toxicity); BIOL (Biological study) (protection from poisoning from; methods and compns. for enhancing cellular function through protection of tissue components such as

muscarinic receptors by administering **pyrophosphate** analogs  
and combination with other agents)

IT 506-32-1, Arachidonic acid 630-08-0, Carbon monoxide, biological studies  
10102-43-9, Nitric oxide, biological studies

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(receptors for, protection of; methods and compns. for enhancing  
cellular function through protection of tissue components such as  
muscarinic receptors by administering **pyrophosphate** analogs  
and combination with other agents)

L117 ANSWER 4 OF 10 HCAPLUS COPYRIGHT 2003 ACS

AN 2001:564841 HCAPLUS

DN 135:132470

TI Selective estrogen receptor modulators in combination with estrogens for  
therapeutic use

IN Labrie, Fernand

PA Endorecherche, Inc., Can.

SO PCT Int. Appl., 160 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K031-565


ICS A61K031-35; A61P015-12; A61P019-10

CC 1-12 (Pharmacology)

Section cross-reference(s): 2, 27, 63

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001054699	A1	20010802	WO 2001-CA86	20010126 <--
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
EP	1251855	A1	20021030	EP 2001-902194	20010126 <--
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
BR	2001008107	A	20030311	BR 2001-8107	20010126 <--
US	2002198179	A1	20021226	US 2001-52803	20011107 <--
US	2003040510	A1	20030227	US 2001-52824	20011107 <--
US	2003065008	A1	20030403	US 2002-143894	20020509 <--
NO	2002003484	A	20020722	NO 2002-3484	20020722 <--
PRAI	US 2000-178601P	P	<del>20000128</del>		<--
	US 2001-771180	A1	20010126		
	WO 2001-CA86	W	20010126		
OS	MARPAT 135:132470				
GI					



- AB Methods for redn. or elimination of the incidence of hot flashes and menopausal symptoms, while decreasing the risk of acquiring breast or endometrial cancer and furthermore treating and/or inhibiting the development of osteoporosis, hypercholesterolemia, hyperlipidemia, atherosclerosis, hypertension, insulin resistance, diabetes, loss of muscle mass, obesity, irregular menstruation, **Alzheimer's** disease, or vaginal dryness in susceptible warm-blooded animals, including humans, involves administration of selective estrogen receptor modulators, particularly compds. I (R1, R2 = OH, moiety convertible to OH in vivo; R3 = (un)satd. (substituted) pyrrolidinyl, (un)satd. (substituted) piperidinyl, etc.) and an amt. of an estrogen or mixed estrogenic/androgenic compd. Further administration of **bisphosphonates**, or a sex steroid precursor is specifically disclosed for the medical treatment and/or inhibition of development of some of these above-mentioned diseases. Pharmaceutical compns. for delivery of active ingredient(s) and kit(s) useful to the invention are also disclosed.
- ST therapeutic estrogen receptor modulator estrogen combination; receptor estrogen modulator prepn therapeutic
- IT Antiarteriosclerotics  
(antiatherosclerotics; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Estrogens  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(antiestrogens; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Mammary gland  
(breast tenderness from hormone replacement therapy; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Drug delivery systems  
(capsules; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Estrogens  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(conjugated; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Bone  
(demineralization; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Menopause  
(disorder, hot flash, and vasomotor symptoms; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Sleep  
(disorder; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Uterus, neoplasm  
(endometrium, inhibitors; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Antitumor agents  
Uterus  
(endometrium; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Uterus  
(epithelium; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Menstrual disorder  
(irregular menstruation; selective estrogen receptor modulators in

- combination with estrogens for therapeutic use)
- IT Muscle
  - (loss of muscle mass; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Antitumor agents
  - (mammary gland; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Mammary gland
  - (neoplasm, inhibitors; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Drug delivery systems
  - (oral; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Estrogens
  - RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
  - (phytoestrogens; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Drug delivery systems
  - (prodrugs; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT **Anti-Alzheimer's agents**
  - Anticholesteremic agents
  - Antidiabetic agents
  - Antihypertensives
  - Antiobesity agents
  - Drug delivery systems
  - Drug interactions
  - Headache
  - Hypolipemic agents
  - Menopause
    - (selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Androgens
  - Estrogens
  - RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
  - (selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Estrogen receptors
  - RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
  - (selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Osteoporosis
  - (therapeutic agents; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Hormone replacement therapy
  - (vaginal bleeding and breast tenderness from; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT Vagina
  - (vaginal bleeding from hormone replacement therapy and vaginal dryness; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT 128607-22-7
  - RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
  - (Fc 1271; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT 130064-21-0P 182167-59-5P 252353-10-9P

- RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(prepn. and reaction; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT 3144-16-9 17720-60-4 151533-33-4  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(reaction; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT 9004-10-8, Insulin, biological studies  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(resistance; selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT 182167-03-9  
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
(selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT 13311-84-7, Flutamide  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
(selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT 50-27-1, Estriol 50-27-1D, Estriol, esters 50-28-2, 17.beta.-Estradiol, biological studies 50-28-2D, 17.beta.-Estradiol, esters 53-16-7, Estrone, biological studies 53-16-7D, Estrone, esters 53-43-0, Dehydroepiandrosterone 53-43-0D, Dehydroepiandrosterone, prodrug derivs. 56-53-1, Diethylstilbestrol 57-63-6, 17.alpha.-Ethinylestradiol 57-63-6D, 17.alpha.-Ethinylestradiol, esters 57-91-0, 17.alpha.-Estradiol 57-91-0D, 17.alpha.-Estradiol, esters 58-22-0, Testosterone 58-22-0D, Testosterone, prodrug derivs. 58-72-0D, Triphenylethylene, derivs. 63-05-8, 4-Androstene-3,17-dione 63-05-8D, 4-Androstene-3,17-dione, prodrug derivs. 72-33-3, Mestranol 72-33-3D, Mestranol, esters 85-95-0, Chemestrogen 120-72-9D, Indole, derivs. 254-04-6D, Benzopyran, derivs. 474-86-2, Equilin 474-86-2D, Equilin, esters 521-17-5, Androst-5-ene-3.beta.,17.beta.-diol 521-17-5D, Androst-5-ene-3.beta.,17.beta.-diol, prodrug derivs. 651-48-9, Dehydroepiandrosterone sulfate 651-48-9D, Dehydroepiandrosterone sulfate, prodrug derivs. 5630-53-5, Tibolone 10540-29-1, Tamoxifen 11095-43-5D, Benzothiophene, derivs. **13598-36-2D, Phosphonic acid, bisphosphonate derivs.** 16005-17-7, Ethynediol 31477-60-8 31477-60-8D, Centchroman, derivs. 68047-06-3, Hydroxytamoxifen 82413-20-5, Droloxifene 84449-90-1, Raloxifene 89778-26-7, Toremifene 116057-75-1, Idoxifene 151751-78-9, 2'-Ethylestrogenoxazole 155701-61-4, GW5638 175737-59-4, LY 326315 180916-16-9, Lasofoxifene 182133-25-1, LY 353381 198481-33-3, TSE 424 252555-01-4 252555-02-5, EM 1520 252555-03-6, EM 1533 318295-61-3, LY 335563 318295-64-6 318295-65-7 318295-66-8 352233-83-1, HMR 3339 352233-84-2, HMR 3656 352233-85-3, LY 335124 352233-86-4, SH 646 352233-89-7, ERA 923 352233-92-2 352233-93-3 352233-94-4  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); **THU (Therapeutic use)**; BIOL (Biological study); USES (Uses)  
(selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- IT 57-88-5, Cholesterol, biological studies 9001-78-9, Alkaline phosphatase  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(selective estrogen receptor modulators in combination with estrogens for therapeutic use)
- RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) American Home Prod; EP 0802183 A 1997 HCAPLUS
- (2) American Home Prod; WO 9909007 A 1999 HCAPLUS
- (3) Behrens, S; WO 0061123 A 2000 HCAPLUS
- (4) Couillard, S; JOURNAL OF THE NATIONAL CANCER INSTITUTE 1998, V90(10), P772 HCAPLUS
- (5) Deshaies, Y; WO 0101969 A 2001 HCAPLUS
- (6) Endorecherche Inc; WO 9963974 A 1999 HCAPLUS
- (7) Labrie, F; US 5776923 A 1998 HCAPLUS
- (8) Labrie, F; JOURNAL OF STEROID BIOCHEMISTRY AND MOLECULAR BIOLOGY 1999, V69(1 - 06), P51

L117 ANSWER 5 OF 10 HCAPLUS COPYRIGHT 2003 ACS

AN 2001:21335 HCAPLUS

DN 134:91140

TI Memory-improving compositions containing astaxanthin or its esters

IN Yamashita, Eiji; Hagino, Nobuyoshi

PA Itano Refrigerated Food Co., Ltd., Japan

SO Jpn. Kokai Tokkyo Koho, 10 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

IC ICM A61K031-12

ICS A23L001-29; A23L002-52; A61K031-00

CC 63-6 (Pharmaceuticals)

Section cross-reference(s): 1, 18

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2001002569	A2	20010109	JP 1999-172757	19990618 <--
PRAI	JP 1999-172757		19990618	<--	
AB	The invention provides a memory-improving compn. contg. astaxanthin or its deriv. as a main component. The compn. is suitable for use in a health food. The effect of Astax-1700 contg. astaxanthin esters and polyunsatd. fatty acid triglycerides, on aging-assocd. memory disorder in mouse was examd. Also, tablets were formulated from Astax-1700 5, microcryst. cellulose 20, magnesium stearate 5 g.				
ST	memory improvement astaxanthin ester				
IT	Drug delivery systems (capsules; memory-improving compns. contg. astaxanthin esters)				
IT	<b>Memory, biological</b> (disorder; memory-improving compns. contg. astaxanthin esters)				
IT	Beverages (health; memory-improving compns. contg. astaxanthin esters)				
IT	Capsules Feed Health food <b>Memory, biological</b> Tablets (memory-improving compns. contg. astaxanthin esters)				
IT	Antioxidants (memory-improving compns. contg. astaxanthin esters and antioxidants)				
IT	Flavonoids Phenols, biological studies RL: FFD (Food or feed use); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (memory-improving compns. contg. astaxanthin esters and antioxidants)				
IT	Phospholipids, biological studies RL: FFD (Food or feed use); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (memory-improving compns. contg. astaxanthin esters and polyunsatd. fatty acid-contg. glycerides or phospholipids)				
IT	Candy				



Chewing gum  
 (memory-improving compns. contg. astaxanthin esters for health foods)

IT Glycerides, biological studies  
 RL: FFD (Food or feed use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (polyunsatd. fatty acid-contg.; memory-improving compns. contg. astaxanthin esters and polyunsatd. fatty acid-contg. glycerides or phospholipids)

IT Fatty acids, biological studies  
 RL: FFD (Food or feed use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (polyunsatd., esters; memory-improving compns. contg. astaxanthin esters and polyunsatd. fatty acid-contg. glycerides or phospholipids)

IT Condiments  
 (rice-seasoning; memory-improving compns. contg. astaxanthin esters for health foods)

IT Drug delivery systems  
 (tablets; memory-improving compns. contg. astaxanthin esters)

IT 472-61-7, Astaxanthin  
 RL: FFD (Food or feed use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (memory-improving compns. contg. astaxanthin)

IT 317830-42-5 318235-68-6, Astax 1700  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); FFD (Food or feed use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (memory-improving compns. contg. astaxanthin esters)

IT 50-81-7, Vitamin C, biological studies 70-18-8, Glutathion, biological studies 83-86-3, Phytic acid 1406-18-4, Vitamin E 7235-40-7, .beta.-Carotene  
 RL: FFD (Food or feed use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (memory-improving compns. contg. astaxanthin esters and antioxidants)

L117 ANSWER 6 OF 10 HCAPLUS COPYRIGHT 2003 ACS

AN 2000:481198 HCAPLUS

DN 133:219815

TI A method for assaying and treating **Alzheimer's** disease involving **amyloid** precursor protein and heparin or divalent cation

IN Masters, Colin Louis; Bush, Ashley Ian; Beyreuther, Konrad

PA The University of Melbourne, Australia

SO Pat. Specif. (Aust.), 51 pp.

CODEN: ALXXAP

DT Patent

LA English

IC ICM G01N033-573

ICS G01N033-577; C12Q001-68; A61K031-70; A61K031-19; A61K031-44; A61K031-16

CC 9-16 (Biochemical Methods)

Section cross-reference(s): 1, 14

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	AU 701954	B2	19990211	AU 1996-50598	19960410 <--
	AU 9650598	A1	19960711		
PRAI	AU 1996-50598		19960410	<--	

AB The present invention relates to a method of assaying for **Alzheimer's** disease in a human by detg. the relative abundance of one or more forms of **amyloid** precursor protein (APP) or the enzyme responsible for said forms in circulatory fluid and to a method for treating the disease by modulating divalent cation and/or heparin interaction with APP. Heparin-Sepharose eluates of **Alzheimer's** disease and control plasma samples were immunoblotted with MAb 22C11 and

the reflectances of the bands at 130, 110, 65 and 42 kDa were measured by computer-assisted image capture anal. The relative amts. of the four APP derivs., as percentages of total lane signal, were detd. in each plasma sample and averaged. The inventors have identified a zinc binding site and a heparin binding site on APP.

- ST **Alzheimer** disease treatment diagnosis **amyloid**  
precursor protein; divalent cation **amyloid** precursor protein  
**Alzheimer**; zinc heparin **amyloid** precursor protein  
**Alzheimer**
- IT Animal cell line  
(PC12, zinc effect on; a method for assaying and treating  
**Alzheimer's** disease involving **amyloid** precursor  
protein and heparin or divalent cation)
- IT **Alzheimer's disease**  
Blood analysis  
(a method for assaying and treating **Alzheimer's** disease  
involving **amyloid** precursor protein and heparin or divalent  
cation)
- IT **Amyloid precursor proteins**  
RL: ADV (Adverse effect, including toxicity); BOC (Biological occurrence);  
BPR (Biological process); BSU (Biological study, unclassified); PUR  
(Purification or recovery); RCT (Reactant); BIOL (Biological study); OCCU  
(Occurrence); PREP (Preparation); PROC (Process); RACT (Reactant or  
reagent)  
(a method for assaying and treating **Alzheimer's** disease  
involving **amyloid** precursor protein and heparin or divalent  
cation)
- IT Blood plasma  
Brain  
(**amyloid** precursor protein purifn. from, of human; a method  
for assaying and treating **Alzheimer's** disease involving  
**amyloid** precursor protein and heparin or divalent cation)
- IT Cations  
(divalent; a method for assaying and treating **Alzheimer's**  
disease involving **amyloid** precursor protein and heparin or  
divalent cation)
- IT Nervous system  
(function of, assessment of; a method for assaying and treating  
**Alzheimer's** disease involving **amyloid** precursor  
protein and heparin or divalent cation)
- IT Immunoassay  
(immunoblotting; a method for assaying and treating **Alzheimer**  
's disease involving **amyloid** precursor protein and heparin or  
divalent cation)
- IT Drug delivery systems  
(oral; a method for assaying and treating **Alzheimer's** disease  
involving **amyloid** precursor protein and heparin or divalent  
cation)
- IT Nerve  
(toxicity; a method for assaying and treating **Alzheimer's**  
disease involving **amyloid** precursor protein and heparin or  
divalent cation)
- IT Cations  
(trivalent; a method for assaying and treating **Alzheimer's**  
disease involving **amyloid** precursor protein and heparin or  
divalent cation)
- IT Biological transport  
(uptake, of zinc; a method for assaying and treating **Alzheimer**  
's disease involving **amyloid** precursor protein and heparin or  
divalent cation)
- IT 9001-92-7, Protease  
RL: ANT (Analyte); ANST (Analytical study)  
(APP-degrading; a method for assaying and treating **Alzheimer**

- 's disease involving **amyloid** precursor protein and heparin or divalent cation)
- IT 7440-66-6, Zinc, biological studies 9005-49-6, Heparin, biological studies 23713-49-7, Zinc ion, biological studies  
RL: ADV (Adverse effect, including toxicity); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(a method for assaying and treating **Alzheimer's** disease involving **amyloid** precursor protein and heparin or divalent cation)
- IT 7446-70-0, Aluminum chloride, biological studies 7646-85-7, Zinc chloride, biological studies 7733-02-0, Zinc sulfate 7758-89-6, Cuprous chloride 7758-94-3, Ferrous chloride  
RL: ADV (Adverse effect, including toxicity); BSU (Biological study, unclassified); BIOL (Biological study)  
(a method for assaying and treating **Alzheimer's** disease involving **amyloid** precursor protein and heparin or divalent cation)
- IT 9005-49-6D, Heparin, conjugates with Sepharose, processes 9012-36-6D, Sepharose, conjugates with heparin  
RL: PEP (Physical, engineering or chemical process); PROC (Process)  
(**amyloid** precursor protein binding to; a method for assaying and treating **Alzheimer's** disease involving **amyloid** precursor protein and heparin or divalent cation)
- IT 286939-08-0  
RL: PEP (Physical, engineering or chemical process); PRP (Properties); PROC (Process)  
(as peptide of **amyloid** precursor protein, zinc ion binding to; a method for assaying and treating **Alzheimer's** disease involving **amyloid** precursor protein and heparin or divalent cation)
- IT 60-00-4, EDTA, biological studies 70-51-9, Desferrioxamine 83-86-3, Phytic acid 83-86-3D, Phytic acid, derivs. 994-36-5, Sodium citrate 115900-75-9 286014-73-1  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(as zinc binding agent; a method for assaying and treating **Alzheimer's** disease involving **amyloid** precursor protein and heparin or divalent cation)
- IT 149658-70-8  
RL: PEP (Physical, engineering or chemical process); PRP (Properties); PROC (Process)  
(in characterization of zinc-binding site of **amyloid** precursor protein; a method for assaying and treating **Alzheimer's** disease involving **amyloid** precursor protein and heparin or divalent cation)
- IT 286014-74-2 286014-75-3  
RL: PRP (Properties)  
(unclaimed sequence; method for assaying and treating **Alzheimer's** disease involving **amyloid** precursor protein and heparin or divalent cation)

L117 ANSWER 7 OF 10 HCAPLUS COPYRIGHT 2003 ACS

AN 1999:449383 HCAPLUS

DN 131:106661

TI Solid products containing calcium and phosphate and methods for the remineralization and prevention of demineralization of teeth

IN Winston, Anthony E.; Usen, Norman

PA Enamelon, Inc., USA

SO PCT Int: Appl., 54 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K007-16

ICS A61K007-18

CC 62-7 (Essential Oils and Cosmetics)

Section cross-reference(s): 63

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9934772	A1	19990715	WO 1998-US24529	19981124 <--
	W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
	AU 9915270	A1	19990726	AU 1999-15270	19981124 <--
PRAI	US 1998-5045		19980109	<--	
	WO 1998-US24529		19981124	<--	
AB	Solid products for remineralizing dental subsurface lesions and/or mineralizing exposed tubules in dentin contain an anionic component composed of at least one phosphate salt and a cationic component composed of at least one calcium salt. The cationic components and the anionic components are mixed in a carrier component and then coated on an insol., solid substrate. Subsurface lesions and/or exposed dentin tubules in a tooth are remineralized by the rapid and simultaneous release of the calcium and phosphate salts into water and/or saliva such that the subsurface lesions and dentin tubules are permeated by the calcium and phosphate ions. The calcium and phosphate ions ppt. as water-insol. calcium phosphate in the subsurface lesions or dentin tubules. The products may be in the form of dental floss, tooth picks, dental tape, dental adhesives, and implants. E.g., wax-coated nylon 6 fibers were passed through a soln. of the polymeric coating and then the calcium and phosphate salts were dusted onto the wet floss. The floss provides excellent cleaning to the interproximal surfaces of the teeth while delivering an effective amt. of calcium and phosphate salts to remineralize the teeth and to combat dental caries.				
ST	calcium phosphate dentifrice tooth mineralization; dental adhesive implant floss tooth mineralization				
IT	Dental materials and appliances				
	(adhesives; dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)				
IT	Saliva				
	(calcium and phosphate release into; dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)				
IT	Tooth				
	(caries, prevention of; dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)				
IT	Dentifrices				
	(dental floss; dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)				
IT	Beeswax				
	Cotton fibers				
	Tooth mineralization				
	Wood				
	Wool				
	(dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)				
IT	Acetate fibers, biological studies				
	Paraffin waxes, biological studies				
	Polyamide fibers, biological studies				
	Polyamides, biological studies				
	Polyester fibers, biological studies				

Polyoxyalkylenes, biological studies  
 Polypropene fibers, biological studies  
 Rayon, biological studies  
 Waxes

RL: BUU (Biological use, unclassified); DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)

- IT Tooth  
 (dentin; dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)
- IT Tooth  
 (disease, demineralization, prevention of; dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)
- IT Stabilizing agents  
 (divalent metal salts; dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)
- IT Ceramics  
 RL: BUU (Biological use, unclassified); DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (fibers; dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)
- IT Dental materials and appliances  
 (implants; dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)
- IT Acrylic fibers, biological studies  
 RL: BUU (Biological use, unclassified); DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (methacrylate-based; dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)
- IT Hydrocarbon waxes, biological studies  
 RL: BUU (Biological use, unclassified); DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (microcryst.; dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)
- IT Dental materials and appliances  
 (tapes; dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)
- IT Dentifrices  
 (tooth picks; dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)
- IT 62-54-4, Calcium acetate 64-19-7, Acetic acid, biological studies  
 77-92-9, biological studies 87-69-4, biological studies 299-28-5, Calcium gluconate 526-95-4, Gluconic acid 814-80-2, Calcium lactate 6915-15-7, Malic acid 7439-95-4D, Magnesium, salts, biological studies 7440-24-6D, Strontium, salts, biological studies 7440-31-5D, Tin, salts, biological studies 7440-66-6D, Zinc, salts, biological studies 7440-70-2D, Calcium, salts, biological studies **7664-38-2**, Phosphoric acid, biological studies **7664-38-2D**, Phosphoric acid, salts, biological studies 7758-23-8, Monocalcium orthophosphate 9000-01-5, Acacia gum 9000-65-1, Gum tragacanth 9003-39-8, Polyvinylpyrrolidone 9004-53-9, Dextrin 9004-64-2, Hydroxypropyl cellulose **9005-25-8**, Starch, biological studies 10043-52-4, Calcium chloride, biological studies 10124-31-9, Ammonium orthophosphate 10124-37-5, Calcium nitrate 16984-48-8, Fluoride, biological studies 25038-54-4, Poly[imino(1-oxo-1,6-hexanediyl)], biological studies 25153-40-6D, Maleic acid-methyl vinyl ether copolymer, alkyl monoesters 25322-68-3 25322-69-4 25609-89-6, Resyn 28-1310 27214-00-2, Calcium glycerophosphate 58748-38-2, Resyn 28-2930 67016-70-0, Amphomer 106392-12-5, Ethylene oxide-propylene oxide block copolymer  
 RL: BUU (Biological use, unclassified); DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)

IT 10103-46-5, Calcium phosphate

RL: FMU (Formation, unclassified); THU (Therapeutic use); BIOL (Biological study); FORM (Formation, nonpreparative); USES (Uses)

(dental solid products contg. calcium and phosphate salts for remineralization and prevention of caries)

RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

(1) Constanz; US 5336264 A 1994 HCAPLUS

(2) Curtis; US 5209251 A 1993

(3) Grabenstetter; US 4083955 A 1978

(4) Raaf; US 4397837 A 1983 HCAPLUS

(5) Tung; US 5460803 A 1995 HCAPLUS

(6) Winston; US 5571502 A 1996 HCAPLUS

(7) Winston; US 5614175 A 1997 HCAPLUS

(8) Winston; US 5645853 A 1997 HCAPLUS

L117 ANSWER 8 OF 10 HCAPLUS COPYRIGHT 2003 ACS

AN 1999:422633 HCAPLUS

DN 131:40560

TI Human chromosome 1p Alzheimer's disease-related gene encodes type 1 **inositol 1,4,5-triphosphate** receptor (IP3R1), and use of IP3R1 and its gene (IP3R1) in diagnosis and/or treatment of Alzheimer's disease

IN Belouchi, Magid; Fillion, Mario; Fortier, Isabel; Robitaille, Yves; Gauvreau, Denis; Ouellette, Gail

PA Algene Biotechnologies, Can.

SO Can. Pat. Appl., 28 pp.

CODEN: CPXXEB

DT Patent

LA English

IC ICM C12N015-12

ICS A61K048-00; G01N033-566; C12Q001-68; C07K014-705

CC 3-3 (Biochemical Genetics)

Section cross-reference(s): 1, 13, 14

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
CA 2203068	AA	19981018	CA 1997-2203068	19970418 <--
PRAI CA 1997-2203068		19970418 <--		

AB This invention discloses that the human chromosome 1p Alzheimer's disease (AD)-related gene encodes a member of the **inositol** phosphate pathway, specifically type 1 **inositol 1,4,5-triphosphate** receptor (IP3R1). The invention further discloses the use of IP3R1 and its gene (IP3R1) for diagnosis and/or treatment of AD. The invention also described the linkage disequil. mapping used to det. regions of the genome implicated in the physiopathol. of AD.

ST human chromosome 1p **inositol triphosphate** receptor  
type 1 gene; Alzheimers disease diagnosis treatment human chromosome 1p IP3R1 gene

IT Gene, animal

RL: BOC (Biological occurrence); BSU (Biological study, unclassified); **THU (Therapeutic use)**; BIOL (Biological study); OCCU (Occurrence); USES (Uses)

(IP3R1; human chromosome 1p Alzheimer's disease-related gene encodes type 1 **inositol 1,4,5-triphosphate** receptor (IP3R1), and use of IP3R1 and its gene (IP3R1) in diagnosis and/or treatment of Alzheimer's disease)

IT Chromosome

(human 1, 1p; human chromosome 1p Alzheimer's disease-related gene encodes type 1 **inositol 1,4,5-triphosphate** receptor (IP3R1), and use of IP3R1 and its gene (IP3R1) in diagnosis and/or

- treatment of Alzheimer's disease)
- IT **Alzheimer's disease**  
(human chromosome 1p Alzheimer's disease-related gene encodes type 1 **inositol 1,4,5-triphosphate** receptor (IP3R1), and use of IP3R1 and its gene (IP3R1) in diagnosis and/or treatment of Alzheimer's disease)
- IT Genetic mapping  
(linkage; human chromosome 1p Alzheimer's disease-related gene encodes type 1 **inositol 1,4,5-triphosphate** receptor (IP3R1), and use of IP3R1 and its gene (IP3R1) in diagnosis and/or treatment of Alzheimer's disease)
- IT Diagnosis  
(mol.; human chromosome 1p Alzheimer's disease-related gene encodes type 1 **inositol 1,4,5-triphosphate** receptor (IP3R1), and use of IP3R1 and its gene (IP3R1) in diagnosis and/or treatment of Alzheimer's disease)
- IT **Inositol 1,4,5-trisphosphate receptors**  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(type 1; human chromosome 1p Alzheimer's disease-related gene encodes type 1 **inositol 1,4,5-triphosphate** receptor (IP3R1), and use of IP3R1 and its gene (IP3R1) in diagnosis and/or treatment of Alzheimer's disease)
- IT 68247-19-8, **Inositol** phosphate  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(pathway; human chromosome 1p Alzheimer's disease-related gene encodes type 1 **inositol 1,4,5-triphosphate** receptor (IP3R1), and use of IP3R1 and its gene (IP3R1) in diagnosis and/or treatment of Alzheimer's disease)

L117 ANSWER 9 OF 10 HCAPLUS COPYRIGHT 2003 ACS

AN 1998:394208 HCAPLUS

DN 129:58812

TI 2-(4-Methoxyphenyl)pyrazolo(4,3-c)quinolin-3-one pharmaceuticals for enhancing cognition

IN Dawson, Gerard Raphael; MacLeod, Angus Murray; Seabrook, Guy Ralph

PA Merck Sharp & Dohme Limited, UK; Dawson, Gerard Raphael; MacLeod, Angus Murray; Seabrook, Guy Ralph

SO PCT Int. Appl., 14 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K031-47

CC 63-6 (Pharmaceuticals)

Section cross-reference(s): 1, 28

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9824435	A1	19980611	WO 1997-GB3232	19971126 <--
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
AU 9851267	A1	19980629	AU 1998-51267	19971126 <--
AU 731533	B2	20010329		
EP 956020	A1	19991117	EP 1997-945942	19971126 <--
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, FI			
JP 2001505218	T2	20010417	JP 1998-525315	19971126 <--
US 6087372	A	20000711	US 1999-308821	19990525 <--
PRAI GB 1996-25398	A	19961206 <--		

WO 1997-GB3232 W 19971126 <--

AB The present invention provides the use of 2-(4-methoxyphenyl)pyrazolo[4,3-c]quinolin-3-one (I) or its salt for enhancing cognition, particularly in **Alzheimer's** disease. Thus, I was prepd. by the reaction of Et 4-chloroquinoline-3-carboxylate with 4-methoxyphenylhydrazine-HCl and converted to its (1S)-(+)-10-camphorsulfonate salt (II). A tablet contained II 20, lactose 120, microcryst. cellulose 40, PVP 5, and Mg stearate 5 mg.

ST methoxyphenylpyrazoloquinolinone pharmaceutical cognition enhancer prepn;  
pyrazoloquinolinone pharmaceutical cognition enhancer prepn

IT **Alzheimer's disease**

**Cognition enhancers**

((methoxyphenyl)pyrazoloquinolinone pharmaceuticals for enhancing cognition)

IT 13720-94-0 19501-58-7

RL: RCT (Reactant); RACT (Reactant or reagent)

((methoxyphenyl)pyrazoloquinolinone pharmaceuticals for enhancing cognition)

IT 77779-50-1P

RL: RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use);  
BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent);  
USES (Uses)

((methoxyphenyl)pyrazoloquinolinone pharmaceuticals for enhancing cognition)

IT 50-21-5DP, Lactic acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt 64-19-7DP, Acetic acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt, biological studies 65-85-0DP, Benzoic acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt, biological studies 75-75-2DP, Methanesulfonic acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt 77-92-9DP, Citric acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt 79-09-4DP, Propionic acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt 87-69-4DP, Tartaric acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt, biological studies 110-15-6DP, Succinic acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt 110-16-7DP, Maleic acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt 526-95-4DP, Gluconic acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt 594-45-6DP, Ethanesulfonic acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt 3144-16-9DP, Camphorsulfonic acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt 6915-15-7DP, Malic acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt 7647-01-0DP, Hydrochloric acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt, biological studies **7664-38-2DP**, Phosphoric acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt, biological studies 7664-93-9DP, Sulfuric acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt, biological studies 10035-10-6DP, Hydrobromic acid, (methoxyphenyl)pyrazoloquinolinone pharmaceutical salt, biological studies 77779-50-1DP, salts  
RL: SPN (Synthetic preparation); **THU (Therapeutic use)**; BIOL (Biological study); PREP (Preparation); USES (Uses)  
((methoxyphenyl)pyrazoloquinolinone pharmaceuticals for enhancing cognition)

RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Anon; PHYSIOLOGY AND BEHAVIOR 1987, V41, P241
- (2) Bennett, D; US 4595684 A 1986 HCAPLUS
- (3) Centre Nat Rech Scient; WO 9221680 A 1992 HCAPLUS
- (4) Ciba Geigy Ag; EP 0022078 A 1981 HCAPLUS

L117 ANSWER 10 OF 10 HCAPLUS COPYRIGHT 2003 ACS

AN 1997:717818 HCAPLUS

DN 127:355343

TI Use of phosphonic acid esters for the treatment of functional disorders of



the brain and depression, and preparation thereof

IN Maurer, Fritz; Schmidt, Bernard; Lensky, Stephan; Van der Staay, Franz-Josef; Fanelli, Richard Joseph; Britelli, David Ross

PA Troponwerke G.m.b.H. & Co. K.-G., Germany; Bayer A.-G.; Maurer, Fritz; Schmidt, Bernard; Lensky, Stephan; Van der Staay, Franz-Josef; Fanelli, Richard Joseph; Britelli, David Ross

SO PCT Int. Appl., 38 pp.  
CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K031-66  
ICS C07F009-40; C07M007-00

CC 1-11 (Pharmacology)  
Section cross-reference(s): 23, 63

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9739756	A1	19971030	WO 1997-US6469	19970417 <--
	W:			AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM	
	RW:			GH, KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG	
	DE 19616471	A1	19971030	DE 1996-19616471	19960425 <--
	AU 9724623	A1	19971112	AU 1997-24623	19970417 <--
	EP 896540	A1	19990217	EP 1997-920431	19970417 <--
	R:			AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, SI, FI, RO	
	CN 1222853	A	19990714	CN 1997-195722	19970417 <--
	BR 9709300	A	19990810	BR 1997-9300	19970417 <--
	JP 2000510455	T2	20000815	JP 1997-538197	19970417 <--
	ZA 9703538	A	19971118	ZA 1997-3538	19970424 <--
	NO 9804964	A	19981222	NO 1998-4964	19981023 <--
	KR 2000010611	A	20000225	KR 1998-708502	19981023 <--
PRAI	DE 1996-19616471	A	19960425	<--	
	WO 1997-US6469	W	19970417	<--	
OS	MARPAT 127:355343				
AB	Phosphonic acid esters are used for the treatment and prevention of functional disorders of the brain and depression. Some of the compds. of the invention are known; others, e.g. di-Me (1-methanesulfonyloxy-2,2,2-trichloroethane)phosphonate, are prepd. Selected compds. were tested in e.g. a Morris maze test and an active avoidance test.				
ST	phosphonic acid ester brain functional disorder; depression phosphonic acid ester prepn				
IT	Mental disorder (affective; phosphonic acid esters for treatment of brain functional disorders and depression, and prepn. thereof)				
IT	Amines, miscellaneous RL: MSC (Miscellaneous) (basic nitrogen compd. acid-binding agents; phosphonic acid esters for treatment of brain functional disorders and depression, and prepn. thereof)				
IT	Mental disorder (cognitive; phosphonic acid esters for treatment of brain functional disorders and depression, and prepn. thereof)				
IT	<b>Cognition</b> (disorder; phosphonic acid esters for treatment of brain functional disorders and depression, and prepn. thereof)				
IT	Acids, uses RL: CAT (Catalyst use); USES (Uses)				

- (inorg.; phosphonic acid esters for treatment of brain functional disorders and depression, and prepn. thereof)
- IT Antidepressants  
Brain, disease  
**Cognition enhancers**  
Drug delivery systems  
**Learning**  
(phosphonic acid esters for treatment of brain functional disorders and depression, and prepn. thereof)
- IT Anhydrides  
Halogen compounds  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(reaction; phosphonic acid esters for treatment of brain functional disorders and depression, and prepn. thereof)
- IT 1554-63-8 10184-66-4 104602-95-1 198561-88-5 198561-89-6  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(isomers; phosphonic acid esters for treatment of brain functional disorders and depression, and prepn. thereof)
- IT 198561-90-9P 198561-91-0P 198562-05-9P  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(phosphonic acid esters for treatment of brain functional disorders and depression, and prepn. thereof)
- IT 52-68-6 4414-11-3 5952-41-0 10184-68-6 **13598-36-2D**,  
Phosphonic acid, esters 61637-95-4 106692-44-8 106692-45-9  
198561-92-1 198561-93-2 198561-94-3 198561-95-4 198561-96-5  
198561-97-6 198561-98-7 198561-99-8 198562-00-4 198562-01-5  
198562-02-6 198562-03-7 198562-04-8 198562-06-0 198562-07-1  
198562-08-2 198562-09-3 198562-10-6 198562-11-7 198562-12-8  
198562-13-9 198562-14-0 198562-15-1 198562-16-2 198562-17-3  
198562-18-4 198562-19-5 198562-20-8 198562-21-9 198562-22-0  
198562-23-1 198562-24-2 198562-25-3 198562-26-4 198562-27-5  
198562-28-6 198562-29-7 198562-30-0 198562-31-1 198562-32-2  
198562-33-3 198562-34-4 198562-35-5 198562-36-6 198562-37-7  
198562-38-8 198562-39-9 198562-40-2 198562-42-4 198562-44-6  
198562-46-8 198562-50-4 198562-51-5 198562-52-6 198562-53-7  
198562-54-8 198562-55-9 198562-56-0 198562-57-1 198562-58-2  
198562-59-3  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); **THU (Therapeutic use)**; BIOL (Biological study); USES (Uses)  
(phosphonic acid esters for treatment of brain functional disorders and depression, and prepn. thereof)
- IT 108-24-7, Acetic anhydride 124-63-0, Methanesulfonic acid chloride  
630-19-3, Pivalic aldehyde 868-85-9, Dimethyl phosphite 1538-75-6,  
Pivalic anhydride  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(reaction; phosphonic acid esters for treatment of brain functional disorders and depression, and prepn. thereof)

=> fil medline

FILE 'MEDLINE' ENTERED AT 09:45:59 ON 04 JUN 2003

FILE LAST UPDATED: 3 JUN 2003 (20030603/UP). FILE COVERS 1958 TO DATE.

On April 13, 2003, MEDLINE was reloaded. See HELP RLOAD for details.

MEDLINE thesauri in the /CN, /CT, and /MN fields incorporate the MeSH 2003 vocabulary. See <http://www.nlm.nih.gov/mesh/changes2003.html>.

for a description on changes.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d all tot

L143 ANSWER 1 OF 12 MEDLINE  
AN 2002470513 MEDLINE  
DN 22217278 PubMed ID: 12231224  
TI Inactivation of the human brain muscarinic acetylcholine receptor by oxidative damage catalyzed by a low molecular weight endogenous inhibitor from **Alzheimer's** brain is prevented by pyrophosphate analogs, bioflavonoids and other antioxidants.  
AU **Fawcett John R**; Bordayo Elizabeth Z; Jackson Kathy; Liu Howard; Peterson Jennifer; Svitak Aleta; **Frey William H 2nd**  
CS The Alzheimer's Research Center, HealthPartners Research Foundation, Regions Hospital, 640 Jackson Street, St. Paul, MN 55101-2595, USA.  
SO BRAIN RESEARCH, (2002 Sep 20) 950 (1-2) 10-20.  
Journal code: 0045503. ISSN: 0006-8993.  
CY Netherlands  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 200212  
ED Entered STN: 20020917  
Last Updated on STN: 20021227  
Entered Medline: 20021223  
AB Oxidative stress has been implicated as a contributing factor to neurodegeneration in **Alzheimer's** disease. An endogenous, low molecular weight (LMW) inhibitor from **Alzheimer's** brain inactivates the human brain muscarinic acetylcholine receptor (mAChR). The inhibitor prevents agonist and antagonist binding to the mAChR as assessed by radioligand binding studies. The LMW endogenous inhibitor, which has components with molecular weights between 100 and 1000 Da, requires dissolved oxygen and glutathione. Prevention of inactivation of the mAChR with peroxidase suggests that the LMW endogenous inhibitor generates peroxide. Heme, previously shown to be present in the LMW endogenous inhibitor, also inactivates the mAChR in the presence of peroxide. Free radical damage to the muscarinic receptor by the endogenous inhibitor can be prevented through the use of naturally occurring antioxidants including bilirubin, biliverdin, carnosol, myricetin and quercetin. In addition, pyrophosphate, **imidodiphosphate, bisphosphonates** and related compounds also protect the muscarinic receptor from free radical damage. Inactivation of the mAChR by the LMW endogenous inhibitor is likely to be a factor in the continual decline of **Alzheimer's** patients, even those taking acetylcholinesterase inhibitors. Natural antioxidants and pyrophosphate analogs may improve the effectiveness of acetylcholinesterase inhibitors and prove useful in the treatment and prevention of **Alzheimer's** disease since the muscarinic acetylcholine receptor is required for memory, and decreased cholinergic function is a critical deficit in **Alzheimer's** disease.  
CT Check Tags: Human; Support, Non-U.S. Gov't  
**Alzheimer Disease: DT, drug therapy**  
**\*Alzheimer Disease: ME, metabolism**  
**\*Antioxidants: PD, pharmacology**  
**Antioxidants: TU, therapeutic use**  
**\*Bioflavonoids: PD, pharmacology**  
**Bioflavonoids: TU, therapeutic use**  
**Brain: DE, drug effects**  
**Brain: ME, metabolism**  
**Catalysis: DE, drug effects**

Diphosphates: CH, chemistry

\*Diphosphates: PD, pharmacology

Diphosphates: TU, therapeutic use

Dose-Response Relationship, Drug

\*Muscarinic Antagonists: ME, metabolism

Muscarinic Antagonists: PD, pharmacology

\*Nerve Tissue Proteins: ME, metabolism

Nerve Tissue Proteins: PD, pharmacology

\*Oxidative Stress: DE, drug effects

Oxidative Stress: PH, physiology

\*Receptors, Muscarinic: ME, metabolism

CN 0 (Antioxidants); 0 (Bioflavonoids); 0 (Diphosphates); 0  
(Muscarinic Antagonists); 0 (Nerve Tissue Proteins); 0 (Receptors,  
Muscarinic); 0 (endogenous modulator protein)

L143 ANSWER 2 OF 12 MEDLINE

AN 1998239725 MEDLINE

DN 98239725 PubMed ID: 9571042

TI Phosphatidylinositol and inositol involvement in Alzheimer amyloid-beta  
fibril growth and arrest.

AU McLaurin J; Franklin T; Chakrabartty A; Fraser P E

CS Centre for Research in Neurodegenerative Diseases, University of Toronto,  
Canada.

SO JOURNAL OF MOLECULAR BIOLOGY, (1998 Apr 24) 278 (1) 183-94.

Journal code: 2985088R. ISSN: 0022-2836.

CY ENGLAND: United Kingdom

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199806

ED Entered STN: 19980708

Last Updated on STN: 19980708

Entered Medline: 19980625

AB A key pathological feature of Alzheimer's disease is the formation and  
accumulation of amyloid fibres. The major component is the 39 to 42  
residue amyloid-beta peptide (Abeta) which is an internal proteolytic  
fragment of the integral membrane amyloid precursor protein. Aggregation  
of Abeta into insoluble amyloid fibres is a nucleation-dependent event  
that may be modulated by the presence of amyloid-associated molecules.  
Fibril formation is also associated with neurotoxicity which may be the  
result of specific Abeta interactions with membrane proteins and/or  
lipids. Using circular dichroism spectroscopy, tyrosine fluorescence  
spectroscopy and electron microscopy, we have examined the binding of  
Abeta peptides 1-40 (Abeta40) and 1-42 (Abeta42) to the glycolipid,  
phosphatidylinositol (PI), and different inositol headgroups. At pH 6.0  
and in the presence of PI vesicles, both Abeta40 and Abeta42 adopted an  
amyloidogenic beta-structure. In contrast, at neutral pH only Abeta42  
folded into a beta-structure in the presence of PI vesicles. To determine  
whether the induction of beta-structure stemmed from interactions with the  
headgroup of PI, the effects of inositol derivatives on Abeta were also  
examined. At pH 7.0, myo-inositol was sufficient to induce beta-structure  
in Abeta42 but had no effect on the conformation of Abeta40. Myo-inositol  
may promote beta-structure as a result of its ability to be both a  
hydrogen-bond donor and acceptor. Mono-, di- and triphosphorylated forms  
of inositol had reduced ability to induce beta-structure in both peptides.  
The results from this study indicate that interaction of Abeta40 and  
Abeta42 with PI acts as a seed for fibril formation while myo-inositol  
stabilizes a soluble Abeta42 micelle.

Copyright 1998 Academic Press Limited.

CT Check Tags: Animal; Support, Non-U.S. Gov't

\*Alzheimer Disease: ME, metabolism

Alzheimer Disease: PA, pathology

Amyloid beta-Protein: AN, analysis

**Amyloid beta-Protein: ME, metabolism**

Cattle

Inositol: CH, chemistry

\*Inositol: ME, metabolism

Inositol 1,4,5-Trisphosphate: ME, metabolism

Inositol Phosphates: ME, metabolism

Peptide Fragments: AN, analysis

Peptide Fragments: ME, metabolism

\*Phosphatidylinositols: ME, metabolism

\*Protein Conformation

Senile Plaques: ME, metabolism

Senile Plaques: PA, pathology

Time Factors

RN 15421-51-9 (inositol 1-phosphate); 47055-78-7 (inositol  
 1,4-bis (phosphate)); 6917-35-7 (Inositol); 85166-31-0 (Inositol  
 1,4,5-Trisphosphate)

CN 0 (Amyloid beta-Protein); 0 (Inositol Phosphates); 0 (Peptide Fragments);  
 0 (Phosphatidylinositols); 0 (amyloid beta-protein (1-40)); 0  
 (beta-amyloid (1-42))

L143 ANSWER 3 OF 12 MEDLINE

AN 1998166903 MEDLINE

DN 98166903 PubMed ID: 9506003

TI Prevention and treatment of osteoporosis: does the future belong to  
 hormone replacement therapy?.

AU Gibaldi M

CS Department of Pharmaceutics, School of Pharmacy, University of Washington,  
 Seattle 98195, USA.

SO JOURNAL OF CLINICAL PHARMACOLOGY, (1997 Dec) 37 (12) 1087-99.

Ref: 76

Journal code: 0366372. ISSN: 0091-2700.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

General Review; (REVIEW)

(REVIEW, TUTORIAL)

LA English

FS Priority Journals

EM 199803

ED Entered STN: 19980410

Last Updated on STN: 19980410

Entered Medline: 19980330

AB Estrogen replacement therapy (ERT) after menopause prevents the  
 development of osteoporosis and reduces the risk of fracture. Other  
 potential benefits are cardioprotection--probably related to the effects  
 of estrogen on lipid profile and fibrinogen levels--and a delay in the  
 onset of Alzheimer's disease and perhaps amelioration of the disease.  
 ERT, however, increases the risk of endometriosis and endometrial cancer  
 unless given with a progestin for at least 10 days per menstrual cycle.  
 It also results in a small but real increase in breast cancer.  
 Alendronate, a **bisphosphonate**, is the first serious competitor  
 of conjugated equine estrogen for the treatment of osteoporosis. Nearing  
 FDA approval are so-called designer estrogens (e.g., raloxifene), which  
 may selectively prevent osteoporosis with little or no effects on  
 endometrial and breast tissue.

CT Check Tags: Female; Human

\*Alendronate: TU, therapeutic use

**Alzheimer Disease: PC, prevention & control**

Calcitonin: TU, therapeutic use

Calcium: TU, therapeutic use

Coronary Disease: PC, prevention &amp; control

**Diphosphonates: TU, therapeutic use**

Endometrial Neoplasms: CI, chemically induced

Endometriosis: CI, chemically induced

\*Estrogen Replacement Therapy  
 Estrogen Replacement Therapy: AE, adverse effects  
 Fluorides, Topical: TU, therapeutic use  
 Forecasting  
 \*Osteoporosis: DT, drug therapy  
 \*Osteoporosis: PC, prevention & control  
 Osteoporosis, Postmenopausal: PC, prevention & control  
 Pulmonary Embolism: CI, chemically induced  
 Sodium Fluoride: TU, therapeutic use  
 Thrombosis: CI, chemically induced  
 Vitamin D: TU, therapeutic use

RN 1406-16-2 (Vitamin D); 66376-36-1 (Alendronate); 7440-70-2 (Calcium);  
 7681-49-4 (Sodium Fluoride); 9007-12-9 (Calcitonin)  
 CN 0 (**Diphosphonates**); 0 (Fluorides, Topical)

L143 ANSWER 4 OF 12 MEDLINE

AN **95303354** MEDLINE

DN **95303354** PubMed ID: **7783950**

TI Preservation of acetylcholine muscarinic M2 receptor G-protein  
 interactions in the neocortex of patients with Alzheimer's disease.

AU Hernandez-Hernandez A; Adem A; Ravid R; Cowburn R F

CS Department of Biochemistry and Molecular Biology, University of Salamanca,  
 Spain.

SO NEUROSCIENCE LETTERS, (1995 Feb 15) 186 (1) 57-60.

Journal code: 7600130. ISSN: 0304-3940.

CY Ireland

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199507

ED Entered STN: 19950726

Last Updated on STN: 20000303

Entered Medline: 19950714

AB The efficacy of acetylcholine muscarinic M2 receptor-G protein coupling  
 was investigated in Alzheimer's disease and control neocortical membranes  
 by measuring the effects of MgCl2 and 5'-guanylylimidodiphosphate  
 (Gpp[NH]p) on high-affinity [3H]oxotremorine-M ([3H]OXO-M) binding. MgCl2  
 gave similar enhancements of [3H]OXO-M binding in Alzheimer's disease and  
 control occipital cortex. In contrast, MgCl2 enhanced [3H]OXO-M binding  
 was significantly higher in Alzheimer's disease superior temporal cortex,  
 compared to controls. MgCl2 enhanced [3H]OXO-M binding in both the  
 occipital and temporal cortices of the Alzheimer's disease cases was  
 reversed to control levels by Gpp[NH]p. It is concluded that the number  
 of high-affinity muscarinic M2 sites is increased in Alzheimer's disease  
 superior temporal, but not occipital, cortex and that M2 sites in both  
 regions maintain an efficient G-protein coupling.

CT Check Tags: Female; Human; In Vitro; Male; Support, Non-U.S. Gov't  
 Aged

\***Alzheimer Disease: ME, metabolism**

\*Cerebral Cortex: ME, metabolism

\*GTP-Binding Proteins: ME, metabolism

**Guanylyl Imidodiphosphate: PD, pharmacology**

Membranes: ME, metabolism

Muscarinic Antagonists

Oxotremorine: ME, metabolism

\*Receptors, Muscarinic: ME, metabolism

RN **34273-04-6 (Guanylyl Imidodiphosphate)**; 70-22-4 (Oxotremorine)

CN 0 (Muscarinic Antagonists); 0 (Receptors, Muscarinic); EC 3.6.1.-  
 (GTP-Binding Proteins)

L143 ANSWER 5 OF 12 MEDLINE

AN **94326289** MEDLINE

DN **94326289** PubMed ID: **7914148**

TI Adenylyl cyclase activity in Alzheimer's disease brain: stimulatory and inhibitory signal transduction pathways are differently affected.

AU Schnecko A; Witte K; Bohl J; Ohm T; Lemmer B

CS Zentrum der Pharmakologie, Johann Wolfgang Goethe-Universitat, Frankfurt, Germany.

SO BRAIN RESEARCH, (1994 May 2) 644 (2) 291-6.  
Journal code: 0045503. ISSN: 0006-8993.

CY Netherlands

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199409

ED Entered STN: 19940914  
Last Updated on STN: 20000303  
Entered Medline: 19940908

AB Adenylyl cyclase (AC) activity was studied in post mortem hippocampus and cerebellum from eight patients with Alzheimer's disease/senile dementia of the Alzheimer type (AD/SDAT) and seven non-demented control patients. AC was stimulated via stimulatory guanine nucleotide binding proteins (Gs) using guanosine triphosphate (GTP) and GppNHp (both  $10(-4)$  M) or directly with either forskolin ( $10(-4)$  M) or  $Mn^{2+}$  ( $10(-2)$  M). Inhibition of AC via Al-receptors was performed with N6-cyclohexyladenosine (CHA) under basal conditions and in the presence of forskolin ( $10(-5)$  M). In both brain regions AC activity was significantly reduced in AD/SDAT when compared to controls. Under basal conditions and after stimulation via Gs mean reduction in hippocampus and cerebellum was 47.7% and 58.2%, respectively. The reduction was less pronounced after direct activation of the AC, amounting to 21.8% in hippocampus and 28.1% in cerebellum. CHA inhibited basal and forskolin-stimulated AC concentration-dependently by about 20% (basal) and 30% (forskolin). Inhibition by CHA was similar in hippocampus and cerebellum and tended to be more pronounced in AD/SDAT than in controls. Since the reduction of AC activity in AD/SDAT is greater after stimulation via Gs than after direct activation of the catalytic subunit, we suggest that both Gs and the catalytic subunit seem to be impaired. The fact that CHA-mediated inhibition of AC is not significantly different in AD/SDAT and controls, indicates that in contrast to Gs-, inhibitory G-proteins (Gi) coupling to AC remains intact in Alzheimer's disease.

CT Check Tags: Female; Human; Male; Support, Non-U.S. Gov't  
Adenosine: AA, analogs & derivatives  
Adenosine: PD, pharmacology  
Adenylate Cyclase: AI, antagonists & inhibitors  
\*Adenylate Cyclase: ME, metabolism  
Aged  
Aged, 80 and over  
\*Alzheimer Disease: EN, enzymology  
Alzheimer Disease: PP, physiopathology  
Brain: DE, drug effects  
Brain: EN, enzymology  
Cerebellum: DE, drug effects  
Cerebellum: EN, enzymology  
Enzyme Activation: DE, drug effects  
Forskolin: PD, pharmacology  
GTP-Binding Proteins: ME, metabolism  
Guanosine Triphosphate: PD, pharmacology  
Guanylyl Imidodiphosphate: PD, pharmacology  
Hippocampus: DE, drug effects  
Hippocampus: EN, enzymology  
Manganese: PD, pharmacology  
Middle Age  
Neurotransmitters: ME, metabolism  
Receptors, Purinergic P1: AI, antagonists & inhibitors  
\*Signal Transduction: PH, physiology

RN 34273-04-6 (Guanylyl Imidodiphosphate); 36396-99-3

(N(6)-cyclohexyladenosine); 58-61-7 (Adenosine); 66428-89-5 (Forskolin);  
 7439-96-5 (Manganese); 86-01-1 (Guanosine Triphosphate)  
 CN 0 (Neurotransmitters); 0 (Receptors, Purinergic P1); EC 3.6.1.-  
 (GTP-Binding Proteins); EC 4.6.1.1 (Adenylate Cyclase)

L143 ANSWER 6 OF 12 MEDLINE

AN 94230615 MEDLINE

DN 94230615 PubMed ID: 7909814

TI Kinesin and tau bind to distinct sites on microtubules.

AU Marya P K; Syed Z; Fraylich P E; Eagles P A

CS Department of Molecular Biology and Biophysics, Randall Institute, King's  
 College, University of London, UK.

SO JOURNAL OF CELL SCIENCE, (1994 Jan) 107 ( Pt 1) 339-44.

Journal code: 0052457. ISSN: 0021-9533.

CY ENGLAND: United Kingdom

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199406

ED Entered STN: 19940620

Last Updated on STN: 19960312

Entered Medline: 19940606

AB We have used a fluorescent derivative of kinesin, AF-kinesin (kinesin  
 conjugated with 5-(iodoacetamido)fluorescein), to investigate the binding  
 site of kinesin on microtubules and to compare this site with that to  
 which tau binds. Microtubules saturated with tau will bind AF-kinesin in  
 the presence of the ATP analogue, 5'-[beta,gamma-imino]triphosphate  
 (AdoPP[NH]P). This shows that there are distinct binding sites for the  
 two proteins. Further evidence comes from digestion studies where  
 taxol-stabilised microtubules were treated with subtilisin, resulting in  
 the cleavage of C-terminal residues from both the alpha- and beta-tubulin  
 subunits. These treated microtubules can no longer bind tau, but are able  
 to bind AF-kinesin in the presence of AdoPP[NH]P. Finally, AF-kinesin  
 will support the gliding of subtilisin-digested microtubules in the  
 presence of ATP at rates comparable to those obtained with non-digested  
 microtubules. These results show directly that the binding site for  
 kinesin is outside the C-terminal region of tubulin that is removed by  
 subtilisin and is distinct from the binding site of tau.

CT Check Tags: Animal; Support, Non-U.S. Gov't

**Adenylyl Imidodiphosphate: PD, pharmacology**

Binding Sites

Brain: ME, metabolism

Cattle

Chromatography, DEAE-Cellulose

Chromatography, High Pressure Liquid

Electrophoresis, Polyacrylamide Gel

Fluoresceins

Kinesin: IP, isolation & purification

\*Kinesin: ME, metabolism

\*Microtubules: ME, metabolism

Microtubules: UL, ultrastructure

Molecular Weight

Paclitaxel: PD, pharmacology

Tubulin: IP, isolation & purification

\*Tubulin: ME, metabolism

**tau Proteins: IP, isolation & purification**

**\*tau Proteins: ME, metabolism**

RN 25612-73-1 (Adenylyl Imidodiphosphate); 33069-62-4 (Paclitaxel);

63368-54-7 (5-iodoacetamidofluorescein)

CN 0 (Fluoresceins); 0 (Tubulin); 0 (tau Proteins); EC 3.6.1.- (Kinesin)

L143 ANSWER 7 OF 12 MEDLINE

AN 92375342 MEDLINE



DN **92375342** PubMed ID: **1508395**  
TI Preservation of Gi-protein inhibited adenylyl cyclase activity in the  
brains of patients with Alzheimer's disease.  
AU Cowburn R F; O'Neill C; Ravid R; Winblad B; Fowler C J  
CS Department of Geriatric Medicine, Karolinska Institute, Huddinge  
University Hospital, Sweden.  
SO NEUROSCIENCE LETTERS, (1992 Jul 6) 141 (1) 16-20.  
Journal code: 7600130. ISSN: 0304-3940.  
CY Netherlands  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199209  
ED Entered STN: 19921009  
Last Updated on STN: 20000303  
Entered Medline: 19920922  
AB The coupling of inhibitory guanine nucleotide binding (Gi) proteins to the  
adenylyl cyclase signal transduction complex was compared in 4 brain  
regions from a series of Alzheimer's disease and matched control subjects  
by measuring the inhibition of membrane enzyme activities in response to  
guanosine 5'-[beta gamma-imido]**diphosphate** (Gpp[NH]p) and  
aluminium fluoride (AlF4-). Basal adenylyl cyclase activities were  
significantly lower in preparations of angular gyrus and frontal and  
temporal cortices, but not cerebellum, from the Alzheimer's disease cases  
compared to controls. Gpp[NH]p and AlF4- gave significant inhibitions of  
adenylyl cyclase activity in all brain regions. The magnitude of these  
inhibitions, when corrected for altered basal activities, were similar for  
the Alzheimer's disease and control cases. These results indicate that  
there is no impairment of Gi-protein mediated inhibition of adenylyl  
cyclase activity in Alzheimer's disease brain.  
CT Check Tags: Human; Support, Non-U.S. Gov't  
\*Adenylate Cyclase: ME, metabolism  
Aluminum: PD, pharmacology  
\***Alzheimer Disease: EN, enzymology**  
\*Brain: EN, enzymology  
Fluorides: PD, pharmacology  
\*GTP-Binding Proteins: PH, physiology  
\***Guanylyl Imidodiphosphate: PD, pharmacology**  
Signal Transduction: PH, physiology  
RN **34273-04-6 (Guanylyl Imidodiphosphate)**; 7429-90-5 (Aluminum);  
7784-18-1 (aluminum fluoride)  
CN 0 (Fluorides); EC 3.6.1.- (GTP-Binding Proteins); EC 4.6.1.1 (Adenylate  
Cyclase)  
L143 ANSWER 8 OF 12 MEDLINE  
AN **92177412** MEDLINE  
DN **92177412** PubMed ID: **1542114**  
TI Metal ion-induced conformational changes of phosphorylated fragments of  
human neurofilament (NF-M) protein.  
AU Hollosi M; Urge L; Perczel A; Kajtar J; Teplan I; Otvos L Jr; Fasman G D  
CS Institute of Organic Chemistry, L. Eotvos University, Budapest, Hungary.  
NC GM45011 (NIGMS)  
SO JOURNAL OF MOLECULAR BIOLOGY, (1992 Feb 5) 223 (3) 673-82.  
Journal code: 2985088R. ISSN: 0022-2836.  
CY ENGLAND: United Kingdom  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199204  
ED Entered STN: 19920424  
Last Updated on STN: 19970203  
Entered Medline: 19920407  
AB The NF-M subunit of human neurofilaments has a C-terminal repeating 13-mer

sequence. The 13-mer (Lys-Ser-Pro-Val-Pro-Lys-Ser-Pro-Val-Glu-Glu-Lys-Gly) (NF-M13) and 17-mer (Glu-Glu-Lys-Gly)-(NF-M13) sequences were synthesized, as were both the mono- and **diphosphorylated** Ser species. Circular dichroism (c.d.) studies and c.d. titrations with Al<sup>3+</sup> and Ca<sup>2+</sup> were performed. The conformation of the phosphorylated and unphosphorylated material was random in water. Deconvolution of the c.d. spectra, in trifluoroethanol, of the untitrated samples yielded a high content of unordered structure, similar to the poly-L-proline II structure. Titration of the phosphorylated species with Al<sup>3+</sup> or Ca<sup>2+</sup> caused a surprising conformational change to occur, yielding a high content of beta-pleated sheet structure. A mechanism of metal binding to the phosphofragments is proposed which may be relevant to the formation of neurofibrillary tangles in Alzheimer's disease.

CT Check Tags: Human; Support, Non-U.S. Gov't; Support, U.S. Gov't, Non-P.H.S.; Support, U.S. Gov't, P.H.S.

\*Aluminum: CH, chemistry

Amino Acid Sequence

Binding Sites

\*Calcium: CH, chemistry

Circular Dichroism

Molecular Sequence Data

\*Neurofilament Proteins: CH, chemistry

Phosphoproteins: CH, chemistry

Phosphorylation

Protein Conformation

RN 111365-29-8 (neurofilament protein M); 7429-90-5 (Aluminum); 7440-70-2 (Calcium)

CN 0 (Neurofilament Proteins); 0 (Phosphoproteins)

L143 ANSWER 9 OF 12 MEDLINE

AN 92168422 MEDLINE

DN 92168422 PubMed ID: 1838799

TI Preservation of 5-hydroxytryptamine<sub>1A</sub> receptor-G protein interactions in the cerebral cortex of patients with Alzheimer's disease.

AU O'Neill C; Cowburn R F; Wiehager B; Alafuzoff I; Winblad B; Fowler C J

CS Alzheimer's Disease Research Centre, Karolinska Institute, Department of Geriatric Medicine, Huddinge University Hospital, Sweden.

SO NEUROSCIENCE LETTERS, (1991 Nov 25) 133 (1) 15-9.

Journal code: 7600130. ISSN: 0304-3940.

CY Netherlands

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199203

ED Entered STN: 19920417

Last Updated on STN: 20000303

Entered Medline: 19920331

AB The coupling of 5-hydroxytryptamine<sub>1A</sub> (5-HT<sub>1A</sub>) receptors to guanine nucleotide binding (G) proteins was investigated in membranes prepared from frontal and parietal cortices of control and Alzheimer's disease brains by characterising the effect of guanosine 5'-[beta gamma-imido] **diphosphate** (Gpp[NH]p) on [3H]8-hydroxy-2-(di-n-propylamino)-tetralin ([3H]8-OH-DPAT) binding parameters. In the absence of guanine nucleotides, [3H]8-OH-DPAT bound to a single high affinity binding site in all membrane types. The number of [3H]8-OH-DPAT binding sites was significantly decreased in the parietal cortex of Alzheimer's disease samples compared with controls, whereas in the frontal cortex the number of binding sites remained unchanged. Gpp[NH]p reduced the [3H]8-OH-DPAT binding affinity and the number of binding sites to the same degree in both regions in control and Alzheimer's disease cases. [3H]8-OH-DPAT binding was inhibited in a concentration dependent manner with an IC<sub>50</sub> value of approximately 1 microM in all cases. These results suggest that the 5-HT<sub>1A</sub> receptor-G protein complex is functionally intact in these

regions in Alzheimer's disease brain.

CT Check Tags: Female; Human; In Vitro; Male; Support, Non-U.S. Gov't  
8-Hydroxy-2-(di-n-propylamino)tetralin  
Aged

\*Alzheimer Disease: ME, metabolism

Cerebral Cortex: DE, drug effects

\*Cerebral Cortex: ME, metabolism

Frontal Lobe: DE, drug effects

Frontal Lobe: ME, metabolism

\*GTP-Binding Proteins: ME, metabolism

Guanine Nucleotides: PD, pharmacology

**Guanylyl Imidodiphosphate: PD, pharmacology**

Histochemistry

Paraffin Embedding

Parietal Lobe: DE, drug effects

Parietal Lobe: ME, metabolism

Receptors, Serotonin: DE, drug effects

\*Receptors, Serotonin: ME, metabolism

Signal Transduction: DE, drug effects

Tetrahydronaphthalenes: ME, metabolism

RN **34273-04-6 (Guanylyl Imidodiphosphate); 78950-78-4**

(8-Hydroxy-2-(di-n-propylamino)tetralin)

CN 0 (Guanine Nucleotides); 0 (Receptors, Serotonin); 0

(Tetrahydronaphthalenes); EC 3.6.1.- (GTP-Binding Proteins)

L143 ANSWER 10 OF 12 MEDLINE

AN **92099092** MEDLINE

DN **92099092** PubMed ID: **1684616**

TI Regional distribution of somatostatin receptor binding and modulation of  
adenylyl cyclase activity in Alzheimer's disease brain.

AU Bergstrom L; Garlind A; Nilsson L; Alafuzoff I; Fowler C J; Winblad B;  
Cowburn R F

CS Alzheimer's Disease Research Group, Karolinska Institute, Department of  
Geriatric Medicine, Huddinge, Sweden.

SO JOURNAL OF THE NEUROLOGICAL SCIENCES, (1991 Oct) 105 (2) 225-33.

Journal code: 0375403. ISSN: 0022-510X.

CY Netherlands

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199202

ED Entered STN: 19920223

Last Updated on STN: 19980206

Entered Medline: 19920204

AB We have previously reported a reduction in the inhibitory effect of  
somatostatin on adenylyl cyclase activity in the superior temporal cortex  
of a group of Alzheimer's disease cases, compared to a group of matched  
controls. In the present study, the levels of high affinity  
125I-Tyr11-somatostatin-14 binding, its modulation by guanine nucleotides  
and the effects of somatostatin on adenylyl cyclase activity have been  
measured in preparations of frontal cortex, hippocampus, caudate nucleus  
and cerebellum from the same patient and control groups. A significant  
reduction in 125I-Tyr11-somatostatin-14 binding was observed in the  
frontal cortex, but not other regions, of the Alzheimer's disease group,  
compared with control values. The profiles of inhibition of specific  
125I-Tyr11-somatostatin-14 binding by Gpp(NH)p were similar in all regions  
in both groups. No significant differences in basal, forskolin-  
stimulated, or somatostatin and neuropeptide Y inhibitions of adenylyl  
cyclase activity were found between the two groups. The pattern of change  
of somatostatin binding in the Alzheimer's disease cases observed in the  
present study differs from the reported pattern of loss of somatostatin  
neurons and may be secondary to the degeneration of somatostatin  
receptor-bearing cholinergic afferents arising from the nucleus basalis.

The results of this study indicate that impaired somatostatin modulation of adenylyl cyclase is not a global phenomenon in Alzheimer's disease brain and also that there are no major disruptions of somatostatin receptor-G-protein coupling or of adenylyl cyclase catalytic activity in this disorder.

CT Check Tags: Human; Support, Non-U.S. Gov't

\*Adenylate Cyclase: ME, metabolism

Aged

Aged, 80 and over

\*Alzheimer Disease: ME, metabolism

\*Brain: ME, metabolism

Caudate Nucleus: ME, metabolism

Cerebellar Cortex: ME, metabolism

Cerebral Cortex: ME, metabolism

Guanylyl Imidodiphosphate: PD, pharmacology

Hippocampus: ME, metabolism

Kinetics

Organ Specificity

Receptors, Neurotransmitter: DE, drug effects

\*Receptors, Neurotransmitter: ME, metabolism

Receptors, Somatostatin

Reference Values

\*Somatostatin: ME, metabolism

RN 34273-04-6 (Guanylyl Imidodiphosphate); 51110-01-1

(Somatostatin)

CN 0 (Receptors, Neurotransmitter); 0 (Receptors, Somatostatin); EC 4.6.1.1 (Adenylate Cyclase)

L143 ANSWER 11 OF 12 MEDLINE

AN 91274844 MEDLINE

DN 91274844 PubMed ID: 2054615

TI Reduced basal and stimulated (isoprenaline, Gpp(NH)p, forskolin) adenylate cyclase activity in Alzheimer's disease correlated with histopathological changes.

AU Ohm T G; Bohl J; Lemmer B

CS Zentrum der Morphologie, J.W. Goethe-Universitat, Frankfurt, Germany.

SO BRAIN RESEARCH, (1991 Feb 1) 540 (1-2) 229-36.

Journal code: 0045503. ISSN: 0006-8993.

CY Netherlands

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199107

ED Entered STN: 19910818

Last Updated on STN: 19980206

Entered Medline: 19910731

AB Cyclic adenosine monophosphate (cAMP) is an adenylate cyclase borne second messenger involved in basic metabolic events. The beta-adrenoceptor sensitive adenylate cyclase was studied in post-mortem hippocampi of controls and Alzheimer patients. Virtually identical subsets of each hippocampus homogenate were stimulated by 100 mumol isoprenaline, Gpp(NH)p and forskolin, respectively, in presence of an ATP-regenerating system. The determination of cAMP formed was carried out by means of a radioassay. The observed significant 50% reduction in basal as well as in stimulated adenylate cyclase activity in Alzheimer's disease is negatively correlated with semiquantitative evaluations of amyloid plaques (P less than 0.05) but not with neuritic plaques, neurofibrillary tangles or neuropil threads. This reduction in enzyme activity is obviously not due to simple cell loss alone. It is likely that the crucial point of the observed functional disturbance is at the level of the catalytic unit of the adenylate cyclase, since the same degree of reduction is maintained at all steps of the signal cascade.

CT Check Tags: Female; Human; Male; Support, Non-U.S. Gov't

\*Adenylate Cyclase: ME, metabolism

Aged

Aged, 80 and over

\*Alzheimer Disease: EN, enzymology

Alzheimer Disease: PA, pathology

\*Brain: EN, enzymology

Brain: PA, pathology

\*Forskolin: PD, pharmacology

\*Guanylyl Imidodiphosphate: PD, pharmacology

\*Isoproterenol: PD, pharmacology

Kinetics

Postmortem Changes

Reference Values

RN 34273-04-6 (Guanylyl Imidodiphosphate); 66428-89-5 (Forskolin);  
7683-59-2 (Isoproterenol)

CN EC 4.6.1.1 (Adenylate Cyclase)

L143 ANSWER 12 OF 12 MEDLINE

AN: 90099495 MEDLINE

DN 90099495 PubMed ID: 2557639

TI Reduced cAMP-signal transduction in postmortem hippocampus of demented old people.

AU Ohm T G; Bohl J; Lemmer B

CS Zentren der Morphologie, J.W. Goethe-Universitat, Frankfurt, FRG.

SO PROGRESS IN CLINICAL AND BIOLOGICAL RESEARCH, (1989) 317 501-9.

Journal code: 7605701. ISSN: 0361-7742.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199001

ED Entered STN: 19900328

Last Updated on STN: 19980206

Entered Medline: 19900131

AB The basal as well as the stimulated activity of the adenylate cyclase was determined in postmortem hippocampi. The tissue probes were obtained from 12 demented individuals (10 Alzheimer-type dementia; 1 Down's syndrome; 1 argyrophilic grains syndrome) and from 15 age-matched controls. The diagnoses were done in accordance with histopathological criteria. Adenylate cyclase was stimulated by isoprenaline, Gpp(NH)p, or forskolin. The amount of cAMP formed was determined by the protein binding method using a radioimmuno assay. In tissues of controls as well as of demented patients adenylate cyclase was stimulated in the rank order of isoprenaline less than Gpp (NH) p less than forskolin. In hippocampal tissues of demented individuals a significant reduction (50%, p less than 0.01) in basal as well as stimulated adenylate cyclase activity was found. This reduction in cAMP signal transduction is not caused by simple cell loss.

CT Check Tags: Female; Human; Male

\*Adenylate Cyclase: ME, metabolism

Aged

Alzheimer Disease: EN, enzymology

\*Alzheimer Disease: ME, metabolism

\*Cyclic AMP: ME, metabolism

Enzyme Activation: DE, drug effects

Forskolin: PD, pharmacology

Guanylyl Imidodiphosphate: PD, pharmacology

Hippocampus: EN, enzymology

\*Hippocampus: ME, metabolism

Isoproterenol: PD, pharmacology

Middle Age

\*Signal Transduction

RN 34273-04-6 (Guanylyl Imidodiphosphate); 60-92-4 (Cyclic AMP);

66428-89-5 (Forskolin); 7683-59-2 (Isoproterenol)  
CN EC 4.6.1.1 (Adenylate Cyclase)

=> fil biosis

FILE 'BIOSIS' ENTERED AT 09:47:31 ON 04 JUN 2003  
COPYRIGHT (C) 2003 BIOLOGICAL ABSTRACTS INC.(R)

FILE COVERS 1969 TO DATE.

CAS REGISTRY NUMBERS AND CHEMICAL NAMES (CNs) PRESENT  
FROM JANUARY 1969 TO DATE.

RECORDS LAST ADDED: 28 May 2003 (20030528/ED)

=> d all tot

L148 ANSWER 1 OF 2 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

AN 2002:589372 BIOSIS

DN PREV200200589372

TI Inactivation of the human brain muscarinic acetylcholine receptor by oxidative damage catalyzed by a low molecular weight endogenous inhibitor from Alzheimer's brain is prevented by pyrophosphate analogs, bioflavonoids and other antioxidants.

AU **Fawcett, John R.**; Bordayo, Elizabeth Z.; Jackson, Kathy; Liu, Howard; Peterson, Jennifer; Svitak, Aleta; **Frey, William H., II** (1)

CS (1) Alzheimer's Research Center, HealthPartners Research Foundation, Regions Hospital, 640 Jackson Street, Saint Paul, MN, 55101-2595: alzheimr@tc.umn.edu USA

SO Brain Research, (20 September, 2002) Vol. 950, No. 1-2, pp. 10-20.  
<http://www.elsevier.com/homepage/sah/bres/doc/journal2.htm>. print.  
ISSN: 0006-8993.

DT Article

LA English

AB Oxidative stress has been implicated as a contributing factor to neurodegeneration in Alzheimer's disease. An endogenous, low molecular weight (LMW) inhibitor from Alzheimer's brain inactivates the human brain muscarinic acetylcholine receptor (mAChR). The inhibitor prevents agonist and antagonist binding to the mAChR as assessed by radioligand binding studies. The LMW endogenous inhibitor, which has components with molecular weights between 100 and 1000 Da, requires dissolved oxygen and glutathione. Prevention of inactivation of the mAChR with peroxidase suggests that the LMW endogenous inhibitor generates peroxide. Heme, previously shown to be present in the LMW endogenous inhibitor, also inactivates the mAChR in the presence of peroxide. Free radical damage to the muscarinic receptor by the endogenous inhibitor can be prevented through the use of naturally occurring antioxidants including bilirubin, biliverdin, carnosol, myricetin and quercetin. In addition, pyrophosphate, imidodiphosphate, bisphosphonates and related compounds also protect the muscarinic receptor from free radical damage. Inactivation of the mAChR by the LMW endogenous inhibitor is likely to be a factor in the continual decline of Alzheimer's patients, even those taking acetylcholinesterase inhibitors. Natural antioxidants and pyrophosphate analogs may improve the effectiveness of acetylcholinesterase inhibitors and prove useful in the treatment and prevention of Alzheimer's disease since the muscarinic acetylcholine receptor is required for memory, and decreased cholinergic function is a critical deficit in Alzheimer's disease.

CC Biochemical Studies - General \*10060

Behavioral Biology - Human Behavior \*07004

Biochemical Studies - Proteins, Peptides and Amino Acids \*10064

Biochemical Studies - Porphyrins and Bile Pigments \*10065

Enzymes - General and Comparative Studies; Coenzymes \*10802  
 Nervous System - Physiology and Biochemistry \*20504  
 Nervous System - Pathology \*20506  
 Psychiatry - Psychopathology; Psychodynamics and Therapy \*21002

BC Hominidae 86215  
 IT Major Concepts  
     Biochemistry and Molecular Biophysics; Nervous System (Neural Coordination)  
 IT Parts, Structures, & Systems of Organisms  
     brain: nervous system  
 IT Diseases  
     Alzheimer's disease: behavioral and mental disorders, nervous system disease  
 IT Chemicals & Biochemicals  
     acetylcholinesterase inhibitors; antioxidants; bilirubin; biliverdin; bioflavonoids; bisphosphonates; carnosol; glutathione; imidodiphosphate; muscarinic acetylcholine receptor: inactivation; muscarinic receptor; myricetin; oxygen; peroxidase; phosphate analogs; pyrophosphate; quericetin  
 IT Alternate Indexing  
     Alzheimer Disease (MeSH)  
 IT Methods & Equipment  
     radioligand binding studies: assessment method, radiobiology method

ORGN Super Taxa  
     Hominidae: Primates, Mammalia, Vertebrata, Chordata, Animalia  
 ORGN Organism Name  
     human (Hominidae)  
 ORGN Organism Superterms  
     Animals; Chordates; Humans; Mammals; Primates; Vertebrates

RN 635-65-4 (BILIRUBIN)  
 114-25-0 (BILIVERDIN)  
 5957-80-2 (CARNOSOL)  
 70-18-8 (GLUTATHIONE)  
 112319-85-4 (IMIDODIPHOSPHATE)  
 529-44-2 (MYRICETIN)  
 7782-44-7 (OXYGEN)  
 9003-99-0 (PEROXIDASE)  
     **14000-31-8 (PYROPHOSPHATE)**

L148 ANSWER 2 OF 2 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.  
 AN 2002:368901 BIOSIS  
 DN PREV200200368901  
 TI Protection of the human brain muscarinic acetylcholine receptor from damage by free radicals generated by an endogenous, low molecular weight inhibitor isolated from Alzheimer's disease brain.  
 AU Bordayo, Elizabeth Z. (1); **Fawcett, John R. (1); Frey, William H., II (1)**  
 CS (1) Alzheimer's Research, HealthPartners Research Foundation, 640 Jackson Street, Saint Paul, MN, 55101 USA  
 SO FASEB Journal, (March 20, 2002) Vol. 16, No. 4, pp. A546.  
<http://www.fasebj.org/>. print.  
 Meeting Info.: Annual Meeting of the Professional Research Scientists on Experimental Biology New Orleans, Louisiana, USA April 20-24, 2002  
 ISSN: 0892-6638.  
 DT Conference  
 LA English  
 AB Current Alzheimer's disease (AD) treatment focuses on the use of cholinesterase inhibitors to increase acetylcholine. This mechanism will only be beneficial if there are viable muscarinic acetylcholine receptors (mAChR) to bind the acetylcholine. In AD brain there is an elevation of a low molecular weight (LMW) inhibitor containing heme that can oxidatively damage the mAChR. This may decrease the efficacy of anticholinesterase therapy. Agonist and antagonist binding to the human brain mAChR was

assessed using radioligand-binding assays. The mAChR was exposed to the LMW inhibitor in the presence and absence of various compounds to determine their effectiveness at protecting the mAChR from oxidative damage. Bile pigments (bilirubin and biliverdin), flavonoids (myricetin and quercetin) and diphosphates (pyrophosphate, imidodiphosphate and bisphosphonate) are hundreds of times more effective than vitamin E at protecting the mAChR. Protecting the human brain mAChR from oxidative stress in aging and AD should help to maintain memory function and increase the effectiveness of cholinesterase inhibitors, cholinergic agonists and related therapeutic agents.

CC General Biology - Symposia, Transactions and Proceedings of Conferences, Congresses, Review Annuals \*00520  
 Behavioral Biology - Human Behavior \*07004  
 Biochemical Studies - General \*10060  
 Nervous System - Physiology and Biochemistry \*20504  
 Nervous System - Pathology \*20506  
 Psychiatry - Psychopathology; Psychodynamics and Therapy \*21002

BC Hominidae 86215

IT Major Concepts  
 Nervous System (Neural Coordination)

IT Diseases  
 Alzheimer's disease: behavioral and mental disorders, nervous system disease

IT Chemicals & Biochemicals  
 bile pigments: free radical-induced muscarinic acetylcholine receptor damage protective effects; diphosphates: free radical-induced muscarinic acetylcholine receptor damage protective effects; endogenous low molecular weight inhibitor protein: Alzheimer disease brain isolation, free radical generation, muscarinic acetylcholine receptor damage inducer; flavonoids: free radical-induced muscarinic acetylcholine receptor damage protective effects

IT Alternate Indexing  
 Alzheimer Disease (MeSH)

IT Miscellaneous Descriptors  
 Meeting Abstract

ORGN Super Taxa  
 Hominidae: Primates, Mammalia, Vertebrata, Chordata, Animalia

ORGN Organism Name  
 human (Hominidae): normal subjects

ORGN Organism Superterms  
 Animals; Chordates; Humans; Mammals; Primates; Vertebrates

RN 14000-31-8 (DIPHOSPHATES)

=> fil wpix

FILE 'WPIX' ENTERED AT 10:18:01 ON 04 JUN 2003

COPYRIGHT (C) 2003 THOMSON DERWENT

FILE LAST UPDATED: 3 JUN 2003 <20030603/UP>

MOST RECENT DERWENT UPDATE: 200335 <200335/DW>

DERWENT WORLD PATENTS INDEX SUBSCRIBER FILE, COVERS 1963 TO DATE

>>> NEW WEEKLY SDI FREQUENCY AVAILABLE --> see NEWS <<<

>>> SLART (Simultaneous Left and Right Truncation) is now available in the /ABEX field. An additional search field /BIX is also provided which comprises both /BI and /ABEX <<<

>>> PATENT IMAGES AVAILABLE FOR PRINT AND DISPLAY <<<

>>> FOR DETAILS OF THE PATENTS COVERED IN CURRENT UPDATES, SEE <http://www.derwent.com/dwpi/updates/dwpicov/index.html> <<<



>>> FOR A COPY OF THE DERWENT WORLD PATENTS INDEX STN USER GUIDE,  
PLEASE VISIT:  
[http://www.stn-international.de/training\\_center/patents/stn\\_guide.pdf](http://www.stn-international.de/training_center/patents/stn_guide.pdf) <<<

>>> FOR INFORMATION ON ALL DERWENT WORLD PATENTS INDEX USER  
GUIDES, PLEASE VISIT:  
[http://www.derwent.com/userguides/dwpi\\_guide.html](http://www.derwent.com/userguides/dwpi_guide.html) <<<

=> d all abeq tech abex tot

L178 ANSWER 1 OF 3 WPIX (C) 2003 THOMSON DERWENT  
AN 2002-082826 [11] WPIX  
DNC C2002-025015  
TI New method for protecting a tissue component in a subject comprises  
administering at least one **pyrophosphate** analog, the subject  
suffers such things as cancer or **Alzheimer's** disease.  
DC B05  
IN CHEN, X; **FAWCETT, J R; FREY, W H; THORNE, R G**  
PA (FAWC-I) FAWCETT J R; (FREY-I) FREY W H; (HEAL-N) **HEALTHPARTNERS RES**  
**FOUND**  
CYC 95  
PI WO 2001082932 A2 20011108 (200211)\* EN 77p A61K031-661 <--  
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
NL OA PT SD SE SL SZ TR TZ UG ZW  
W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CR CU CZ DE DK DM  
DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC  
LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD SE  
SG SI SK SL TJ TM TR TT TZ UA UG UZ VN YU ZA ZW  
US 2002028786 A1 20020307 (200221) A61K031-7105 <--  
AU 2001057444 A 20011112 (200223) A61K031-661 <--  
EP 1278525 A2 20030129 (200310) EN A61K031-661 <--  
R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
RO SE SI TR  
ADT WO 2001082932 A2 WO 2001-US13931 20010430; US 2002028786 A1 Provisional US  
2000-200843P 20000501, Provisional US 2000-230263P 20000906, Provisional  
US 2000-233025P 20000915, US 2001-844450 20010427; AU 2001057444 A AU  
2001-57444 20010430; EP 1278525 A2 EP 2001-930957 20010430, WO  
2001-US13931 20010430  
FDT AU 2001057444 A Based on WO 200182932; EP 1278525 A2 Based on WO 200182932  
PRAI US 2000-233025P 20000915; US 2000-200843P 20000501; US 2000-230263P  
20000906; US 2001-844450 20010427  
IC ICM **A61K031-661; A61K031-7105**  
ICS **A61K031-66; A61K031-6615; A61K031-662;**  
**A61K031-706; A61K031-7076; A61K031-7084**  
AB WO 200182932 A UPAB: 20020215  
NOVELTY - A method of protecting a tissue component in a subject comprises  
administering at least one **pyrophosphate** analog.  
DETAILED DESCRIPTION - A method of protecting a tissue component in a  
subject comprises administering at least one **pyrophosphate**  
analog of formula (I) or (II) or a dinucleoside-5'-5'-**pyrophosphate**  
, a **cyclopyrophosphate** of purine, a pyrimidine acyclonucleoside,  
an inositol diphosphate, an inositol triphosphate, an inositol  
tetraphosphate, an inositol pentaphosphate, an inositol hexaphosphate or  
their salt.  
X = O, CH<sub>2</sub>, NH or S;  
R<sub>1</sub> = H, lower alkyl, guanyl, adenylyl, glycerol, acyl glycerol,  
diacyl glycerol, serine, threonine, tyrosine, arachidonyl, -PO(OH)(OR<sub>2</sub>),  
or -(PO(OH)O)<sub>m</sub>-PO(OH)(OR<sub>2</sub>);  
m = 1-3;  
R<sub>2</sub> = H, lower alkyl, guanyl, adenylyl, glycerol, acyl glycerol,  
diacyl glycerol, serine, threonine, tyrosine, or arachidonyl; and  
n = 1-900.  
p = 2-4;

Y = O, RCR3, CR, C (p=4), CH (p=3), or CH2 (p=2), NH, N, S;  
 R, R3 = H, OH, lower alkyl, or (CH2)<sub>q</sub>NH2; and  
 q = 1-6.

INDEPENDENT CLAIMS are included for:

- (1) a method of protecting tissue from oxidative stress;
- (2) a method of increasing the efficacy of an agent that affects a receptor;
- (3) a method of protecting a subject from at least one carcinogenic metal;
- (4) a method of reducing poisoning of a subject by at least one metal;
- (5) a method (e) of treating bacterial, fungal, algo, or algae infections;
- (6) a method of reducing toxic actions of metal ions;
- (7) a method (g) of protecting a pharmacological agent in a formulation; and
- (8) a method (h) of increasing efficacy of a neurologic agent.

ACTIVITY - Cytostatic; cardiant; hypotensive; vasotropic; antiinflammatory; uropathic; antidiabetic; immunosuppressive; osteopathic; nootropic; neuroprotective; cerebroprotective; tranquilizer; neuroleptic; antitumor.

MECHANISM OF ACTION - Muscarinic acetylcholine receptor modulator.

USE - The methods are used on subjects suffering from cancer, neuropathies, diseases or disorders of the heart, smooth muscles, blood, blood vessels, glands, or bones, hypertension, myocardial infarction, ischemic heart disease or congestive heart failure, irritable bowel syndrome, diverticular disease, urinary incontinence, esophageal achalasia, chronic obstructive airways disease, cardiac arrhythmia, xerostomia, diabetes mellitus, Sjogren's syndrome, Paget's disease, hereditary hematochromatosis, a non-CNS disorder resulting from normal aging, a neurologic disorder, a psychiatric disorder, **Alzheimer's** disease, Parkinson's disease, Lewy body dementia, multiple sclerosis, cerebellar ataxia, progressive supranuclear palsy, amyotrophic lateral sclerosis, an affective disorder, an anxiety disorder, schizophrenia, cell damage, nerve damage, a CNS infection, a tumor of the brain, a tumor of the spinal cord, a stroke in the brain, a stroke in the spinal cord, a prion disease, a CNS disorder resulting from ordinary aging, a brain injury, a spinal cord injury, or a non-CNS disorder resulting from normal aging.

Dwg.0/20

FS CPI

FA AB; GI; DCN

MC CPI: **B05-B01G; B05-B01J; B05-B01P; B06-D09;**

B07-A02A; B10-A17; B10-B02E; B10-B02H; B10-E04C; B14-E10C; B14-F01;

B14-F01B; B14-F01E; B14-F02; B14-F02B; B14-G02; B14-H01; B14-H01B;

B14-J01; B14-J01A3; **B14-J01A4;** B14-J01B3; B14-J01B4;

B14-J05; B14-L01; B14-L06; B14-N01; B14-N16; B14-S01; B14-S04

TECH UPTX: 20020215

TECHNOLOGY FOCUS - BIOLOGY - Preferred tissue: The tissue component is at least one of receptor, a protein, a lipid, a nucleic acid, a carbohydrate, a hormone, a vitamin and a cofactor.

Preferred receptor: The receptor is for a neurotransmitter, a neuropeptide, a neurotrophin, a growth factor, a steroid, a histamine, a purine, a benzodiazepine, arachidonic acid, nitric oxide, carbon monoxide, an odorant, or an ion channel. When the method is for increasing the efficacy of an agent that affects a receptor, the receptor is one of muscarinic acetylcholine, nicotinic acetylcholine, an opiate, a catecholamine, serotonin, glutamate, aspartate, cannabinoid, gamma aminobutyric acid, or glycine.

Preferred agent: The agent affects a muscarinic acetylcholine receptor and comprises an anticholinesterase agent, a neurologic agent, a muscarinic receptor agonist, xanomeline, donepezil, rivastigmine, galanthamine, metrifonate. The pharmacological agent in method (g) is a therapeutic or

diagnostic agent. The neurologic agent comprises a ganglioside (comprising GM-1 ganglioside), a phosphatidylserine, a nerve growth factor, a neurotrophin (comprising neurotrophin 3, 4 and/or 5), a brain-derived neurotrophic factor, a fibroblast growth factor (comprising basic fibroblast growth factor or acidic fibroblast growth factor), an insulin, an insulin-like growth factor (comprising insulin-like growth factor-I and/or 2), a transforming growth factor, an epidermal growth factor, a platelet-derived growth factor, a neurokinin, activity-dependent neurotrophic factor, a ciliary neurotrophic factor, a glia-derived neurotrophic factor, a glia-derived nexin, a cholinergic enhancing factor (which comprises ethanolamine, thyroid hormone T. 3 and/or gallamine), an antisense oligonucleotide, a DNA or RNA vector or plasmid that encodes one or more protein neurologic agents and/or nerve growth promoting factors. Preferred method: The subject in method (e) is a plant, an animal or a mammal.

TECHNOLOGY FOCUS - INORGANIC CHEMISTRY - Preferred metal: The carcinogenic metal is arsenic, cadmium, cobalt, nickel, lead or chromium. The metal is iron, copper, mercury, lead, cadmium, vanadium or their alloys. The metal ions are Fe<sup>++</sup>, Hg<sup>++</sup>, Cd<sup>++</sup>, Cu<sup>++</sup>, As<sup>++</sup> or Pb<sup>++</sup>.

TECHNOLOGY FOCUS - ORGANIC CHEMISTRY - Preferred method: The methods further comprise combining any **pyrophosphate** analog with bilirubin, biliverdin, carnosol, quercetin, myricetin, bioflavonoid, heme oxygenase, a vector encoding a heme oxygenase, heme oxygenase-1, a vector encoding a heme oxygenase-1, heme oxygenase-2, a vector encoding a biliverdin reductase, a catalase, a vector encoding a catalase, a peroxidase, a vector encoding a peroxidase, and/or a heme binding protein (which comprises hemopexin and/or a lipoprotein).

ABEX

UPTX: 20020215

SPECIFIC COMPOUNDS - The **pyrophosphate** in method (e) comprises imidodiphosphate. Preferably the **pyrophosphate** analog comprises imidodiphosphate, guanylimidodiphosphate, adenylylimidodiphosphate, etidronic acid, and/or pamidronic acid.

L178 ANSWER 2 OF 3 WPIX (C) 2003 THOMSON DERWENT

AN 2000-431196 [37] WPIX

DNC C2000-130995

TI Administration of agent to the central nervous system (CNS) via tissue innervated by the trigeminal nerve, used in the treatment and diagnosis of diseases and disorders of the CNS, brain and spinal cord..

DC A14 A17 A96 B07

IN CHEN, X; FREY, W H; THORNE, R G

PA (CHIR) CHIRON CORP

CYC 90

PI WO 2000033814 A2 20000615 (200037)\* EN 54p A61K009-00

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
OA PT SD SE SL SZ TZ UG ZW

W: AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CR CU CZ DE DK DM EE ES  
FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS  
LT LU LV MA MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL  
TJ TM TR TT TZ UA UG UZ VN YU ZA ZW

AU 2000021734 A 20000626 (200045) A61K009-00

EP 1135105 A2 20010926 (200157) EN A61K009-00

R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
RO SE SI

JP 2002531490 W 20020924 (200278) 72p A61K009-08

ADT WO 2000033814 A2 WO 1999-US29335 19991209; AU 2000021734 A AU 2000-21734  
19991209; EP 1135105 A2 EP 1999-966114 19991209, WO 1999-US29335 19991209;  
JP 2002531490 W WO 1999-US29335 19991209, JP 2000-586308 19991209

FDT AU 2000021734 A Based on WO 200033814; EP 1135105 A2 Based on WO  
200033814; JP 2002531490 W Based on WO 200033814

PRAI US 1998-208539 19981209

IC ICM A61K009-00; A61K009-08  
 ICS A61K009-06; A61K009-107; A61K009-12; A61K009-127; A61K038-00;  
 A61K038-28; A61K045-00; A61K047-24; A61K047-32; A61P025-00;  
 A61P025-08; A61P025-18; A61P025-22; A61P025-24; A61P025-28;  
 A61P027-16; A61P031-18; A61P031-22; A61P035-00

AB WO 200033814 A UPAB: 20000807  
 NOVELTY - A method of delivering agents to the central nervous system (CNS) comprises administering a composition comprising the agent to a tissue innervated by the trigeminal nerve and outside the nasal cavity, wherein the agent is absorbed through the tissue and is transported to the CNS.  
 USE - The method can be used to deliver agents to the CNS including the spine and the brain for diagnosis, treatment or prevention of disorders or diseases of the CNS, brain or spinal cord. The disorders may be neurologic or psychiatric and include **Alzheimer's** disease, Parkinson's disease, Lewy body dementia, multiple sclerosis, epilepsy, cerebellar ataxia, progressive supranuclear palsy, amyotrophic lateral sclerosis, affective disorders, anxiety disorders, obsessive compulsive disorders, personality disorders, attention deficit disorders, attention deficit hyperactivity disorder, Tourette syndrome, Tay Sachs, Nieman Pick and other lipid storage and genetic brain diseases and schizophrenia. The method can also be used in patients suffering from or at risk of nerve damage from cerebrovascular disorders such as stroke, from CNS infections including meningitis and human immunodeficiency virus (HIV), from tumors of the brain and spinal cord or from prion disease, to counter CNS disorders resulting from aging, from brain injury or from spinal cord injury or to treat neurodegenerative disorders.  
 ADVANTAGE - The method is non-invasive, unlike most methods for administration to the CNS.  
 Dwg.0/0

FS CPI  
 FA AB; DCN  
 MC CPI: A04-G07; A12-V01; A12-V03C2; B04-B01B; B04-B03C; B04-C01; B04-C03;  
 B04-D01; B04-E01; B04-E08; B04-H06; B04-H06G; **B05-B01P**;  
 B12-M05; B14-A02B1; B14-H01; B14-J01; B14-J07; B14-N16; B14-S01

TECH UPTX: 20000807  
 TECHNOLOGY FOCUS - PHARMACEUTICALS - Preferred Tissue: The agent is administered to tissue innervated by the trigeminal nerve but outside the nasal cavity, therefore it may be administered to oral tissue e.g. teeth, the gums, the floor of the oral cavity, the cheeks, the lips, the tongue or a combination thereof; to mucosa or epithelium innervated by the trigeminal nerve e.g. mucosa or epithelium of or surrounding the eye, such as mucosa or epithelium of the upper eyelid, the lower eyelid, the eyeballs, the conjunctiva, the lacrimal system or a combination thereof; to the skin of the face, scalp or temporal region suitable skin of the face includes skin of the chin, the upper lip, the forehead, the nose, the cheek, skin around the eyes and combinations thereof; or to tissue of or around the ear e.g. the auricle, the external acoustic meatus, the tympanic membrane, skin in the temporal region especially skin of the temple and the lateral part of the scalp and combinations thereof.  
 Preferred Agents: Agents that can be delivered to the CNS include organic pharmaceuticals, inorganic molecules, peptides, peptoids, proteins, lipids, carbohydrates, nucleic acids or diagnostic agents. Preferred neurologic agents include GM1 ganglioside, fibroblast growth factor especially basic fibroblast growth factor, insulin-like growth factor especially insulin-like growth factor 1, phosphatidyl serine, a plasmid, a vector, an antisense oligonucleotide etc. Preferred Compositions: The composition may comprise a liquid, powder, spray, gel, ointment, infusion or a combination thereof. The composition comprises the active agent and a carrier, additive and/or adjuvant. Among the optional substances in the composition lipophilic substances that can enhance absorption of the agent are particularly preferred. The agent may be mixed with a lipophilic adjuvant alone or in combination with a carrier or may be combined with

one of several types of micelle or liposome substances. Lipophilic micelles and liposomes preferably comprise a ganglioside, a phosphatidylcholine, a phosphatidylserine, lipofectin, DOTAP or a combination thereof. The composition may additionally contain a controlled release polymer comprising a poly(ethylene-co-vinylacetate) especially for sub-lingual administration.

Preferred Route: The agent is transported along a neural pathway which comprises lymphatic channels running with a nerve and is transported to the CNS including to a hippocampal formation, an amygdaloid nuclei, a nucleus basalis of Meynert, a locus ceruleus, a brainstem raphe nuclei or combinations thereof, the spinal cord, the brain stem, a cortical structure, a subcortical structure and any combinations thereof.

ABEX

UPTX: 20000807

ADMINISTRATION - Administration may be continuous or intermittent. Dosage is dependent on a wide variety of factors.

EXAMPLE - Male Sprague-Dawley rats, 200-310g, were anesthetized with intraperitoneal pentobarbital (40 mg/kg). Drug delivery to the brain and spinal cord was assessed after sublingual administration of 7.4 nmol of 125I-IGF-I (insulin-like growth factor I) in phosphate buffered saline, pH 7.4. Rats were placed in on their bellies with posterior elevated and mouth lowered. 125I-IGF-I on a small strip of filter paper was placed under the tongue. The rats subsequently underwent perfusion-fixation within minutes following completion of administration. Areas dissected included selected brain regions and the cervical, thoracic and sacral regions of the spinal cord. Rapid appearance of neurologic agent in the brain and spinal cord was observed by determining the radioactivity. The concentration of the neurologic agent was higher in the cervical region of the spinal cord than in the thoracic region, and was higher in the thoracic region than in the sacral or lumbar regions. High concentrations were found in the meninges or dura surrounding each of the following: the olfactory bulb, the dorsal and ventral regions of the brain, the trigeminal nerve and the upper cervical spinal cord. The IGF was also found in the olfactory bulb, spinal cord and the brain stem. The trigeminal nerve itself contained high concentrations of the neurologic agent.

L178 ANSWER 3 OF 3 WPIX (C) 2003 THOMSON DERWENT

AN 2000-431195 [37] WPIX

DNC C2000-130994

TI Administration of a neurotrophic agent to the central nervous system (CNS) via administration to the nasal cavity used in the treatment and diagnosis of disorders and diseases of the CNS, brain and spinal cord..

DC A14 A17 A96 B07

IN CHEN, X; FREY, W H; THORNE, R G

PA (CHIR) CHIRON CORP

CYC 90

PI WO 2000033813 A1 20000615 (200037)\* EN 62p A61K009-00

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
OA PT SD SE SL SZ TZ UG ZW

W: AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CR CU CZ DE DK DM EE ES  
FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS  
LT LU LV MA MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL  
TJ TM TR TT TZ UA UG UZ VN YU ZA ZW

AU 2000020495 A 20000626 (200045) A61K009-00

EP 1137401 A1 20011004 (200158) EN A61K009-00

R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MK RO SI

JP 2002531489 W 20020924 (200278) 74p A61K045-00

ADT WO 2000033813 A1 WO 1999-US29334 19991209; AU 2000020495 A AU 2000-20495  
19991209; EP 1137401 A1 EP 1999-964208 19991209, WO 1999-US29334 19991209;  
JP 2002531489 W WO 1999-US29334 19991209, JP 2000-586307 19991209

FDT AU 2000020495 A Based on WO 200033813; EP 1137401 A1 Based on WO  
200033813; JP 2002531489 W Based on WO 200033813

PRAI US 1998-208538 19981209

IC ICM A61K009-00; A61K045-00

ICS A61K009-06; A61K009-08; A61K009-12; A61K009-127; A61K009-14;  
A61K009-72; A61K038-00; A61K047-24; A61K047-30; A61K047-32;  
A61K047-36; A61P003-02; A61P025-00; A61P025-08; A61P025-16;  
A61P025-18; A61P025-22; A61P025-28

AB WO 200033813 A UPAB: 20000807

NOVELTY - Method for transporting a neurotrophic agent to the central nervous system (CNS) comprises administering the agent to the upper third of the nasal cavity, wherein the neurotrophic agent is absorbed through the nasal cavity and transported to the CNS.

USE - The method can be used to deliver agents to the CNS including the spine and the brain for diagnosis, treatment or prevention of disorders or diseases of the CNS, brain or spinal cord. The disorders may be neurologic or psychiatric and include **Alzheimer's** disease, Parkinson's disease, Lewy body dementia, multiple sclerosis, epilepsy, cerebellar ataxia, progressive supranuclear palsy, amyotrophic lateral sclerosis, affective disorders, anxiety disorders, obsessive compulsive disorders, personality disorders, attention deficit disorders, attention deficit hyperactivity disorder, Tourette syndrome, Tay Sachs, Nieman Pick and other lipid storage and genetic brain diseases and schizophrenia. The method can also be used in patients suffering from or at risk of nerve damage from cerebrovascular disorders such as stroke, from CNS infections including meningitis and HIV, from tumors of the brain and spinal cord or from prion disease, to counter CNS disorders resulting from aging, from brain injury or from spinal cord injury or to treat neurodegenerative disorders.

ADVANTAGE - The method is non-invasive, unlike most methods for administration to the CNS.

Dwg.0/5

FS CPI

FA AB; DCN

MC CPI: A04-G07; A12-V01; A12-V03C2; B04-B01B; B04-C03B; B04-H06; B04-H06D;  
B04-H06G; **B05-B01P**; B12-M01B; B14-A02B1; B14-J01; B14-N16

TECH UPTX: 20000807

TECHNOLOGY FOCUS - PHARMACEUTICALS - Preferred Administration: The agent is administered to tissue innervated by the trigeminal and olfactory nerves inside the nasal cavity and sinuses. The agent is delivered to the olfactory area in the upper third of the nasal cavity and particularly to the olfactory epithelium in the roof of the nose.

Preferred Agents: Neurotrophic agents that can be delivered to the CNS include fibroblast growth factor especially basic fibroblast growth factor, insulin-like growth factor especially insulin-like growth factor 1 and nerve growth factor.

Preferred Compositions: The composition may comprise a liquid, powder, spray, gel, ointment, infusion or a combination thereof. The composition comprises the active agent and a carrier, additive and/or adjuvant. Among the optional substances in the composition lipophilic substances that can enhance absorption of the agent are particularly preferred. The agent may be mixed with a lipophilic adjuvant alone or in combination with a carrier or may be combined with one of several types of micelle or liposome substances. Lipophilic micelles and liposomes preferably comprise a ganglioside, a phosphatidylcholine, a phosphatidylserine, lipofectin, DOTAP or a combination thereof. The composition may additionally contain a controlled release polymer comprising a poly(ethylene-co-vinylacetate). Preferred Route: The agent is transported along a neural pathway, either trigeminal or olfactory nerve pathway, which comprises lymphatic channels running with a nerve and is transported to the CNS including to an olfactory bulb, a hippocampal formation, a frontal cortex, a midbrain, a brainstem, a spinal cord or a combination thereof.

TECHNOLOGY FOCUS - POLYMERS - Preferred Composition: The composition for administration to the CNS may comprise neurologic agents including fibroblast growth factor especially basic fibroblast growth factor,

insulin-like growth factor especially insulin-like growth factor 1 and nerve growth factor. The composition may comprise a controlled release polymer comprising a poly(ethylene-co-vinylacetate). The composition preferably also comprises a lipophilic adjuvant alone or in combination with a carrier or may be combined with one of several types of micelle or liposome substance.

ABEX UPTX: 20000807

ADMINISTRATION - Administration may be continuous or intermittent. Dosage is approximately 0.1-10 nmol of the neurotrophic agent, resulting in a therapeutically effective amount of 10<sup>-11</sup>M to about 10<sup>-9</sup> M in a portion of the CNS.

EXAMPLE - Male Sprague-Dawley rats, 200-310 g, were anesthetized with intraperitoneal pentobarbital (40 mg/kg). Drug delivery to the brain and spinal cord was assessed after intranasal administration of 7.4 nmol of 125I-IGF-I (insulin-like growth factor I) in phosphate buffered saline, pH 7.4. Rats were placed on their backs and administered about 25 microlitres of 125I-IGF-I to each naris over 10-22 minutes, alternating drops every 2-3 minutes between the left and right nares. The rats subsequently underwent perfusion-fixation within minutes following completion of administration. Areas dissected included the olfactory bulbs, medulla, pons and cerebellum. Rapid appearance of radiolabel in the brain observed with the highest concentrations seen in the olfactory bulbs (3 + 0.47 nM), medulla (0.62 + 0.16 nM), pons (0.31 + 0.07 nM) and cerebellum (0.3 + 0.1 nM).

=> d his

(FILE 'HOME' ENTERED AT 08:06:33 ON 04 JUN 2003)  
SET COST OFF

FILE 'HCAPLUS' ENTERED AT 08:06:47 ON 04 JUN 2003

	E US20020028786/PN
L1	1 S E3
	E WO2001-US13931/AP, PRN
L2	1 S E3, E4
	E US2000-233263/AP, PRN
L3	1 S E5
	E US2000-233025/AP, PRN
L4	1 S E5
	E US2000-230263/AP, PRN
L5	1 S E5
L6	1 S L1-L5
	E HEALTHPARTNER/PA, CS
L7	12 S E5-E11
	E HEALTH PARTNER/PA, CS
L8	23 S (HEALTH(L) PARTN?)/PA, CS
	SEL DN AN 23
L9	1 S L8 AND E1-E3
L10	13 S L7, L9
	E FREY W/AU
L11	167 S E3, E7-E9, E25, E29-E33
	E FAWCETT J/AU
L12	33 S E3, E13
	E FAWCETT JOHN/AU
L13	171 S E3, E6, E7
L14	48821 S ?PYROPHOS?
L15	144 S ?PYRO PHOS?
	E PYROPHOS/CT
	E E16+ALL
L16	851 S E1
	E E2+ALL

L17 8258 S E10+NT  
       E E8+ALL  
 L18 24848 S E4+NT  
       E E3+ALL  
 L19 219974 S E3+NT  
 L20 2 S L10 AND L14-L19  
 L21 3 S L11-L13 AND L14-L19  
 L22 3 S L6,L20,L21  
 L23 2 S L22 NOT RHODIUM/TI  
       E ALZHEIMER/CT  
 L24 11589 S E3-E20  
       E E9+ALL  
 L25 11603 S E6,E5+NT  
 L26 11213 S E23+NT OR E24+NT OR E25+NT OR E26+NT OR E27+NT OR E28+NT OR E  
       E E23+ALL  
 L27 1188 S E8  
       E E6+ALL  
 L28 72281 S E7,E6+NT  
       E E22+ALL  
       E E5+ALL  
       E E25+ALL  
 L29 10751 S E9+NT  
       E E8+ALL  
 L30 2080 S E6,E5+NT  
 L31 28838 S E15+NT OR E13+NT  
       E E10+ALL  
       E E27+ALL  
 L32 449 S E8-E11  
       E E14+ALL  
       E E28+ALL  
       E E3+ALL  
       E E29+ALL  
 L33 961 S E3  
 L34 3 S L10 AND L24-L33  
 L35 14 S L11-L13 AND L24-L33  
 L36 14 S L23,L34,L35  
 L37 4 S L36 AND ?PHOSPH?  
 L38 2 S L37 AND L23  
       SEL RN

FILE 'REGISTRY' ENTERED AT 08:21:28 ON 04 JUN 2003

L39 66 S E1-E67  
 L40 14 S L39 AND P/ELS  
 L41 3 S L40 AND (C4H8CL3O4P OR C2H8O7P2 OR C3H11NO7P2)  
 L42 1 S L40 AND H3O3P  
 L43 2 S L41 AND P>=2  
 L44 10 S L40 NOT L41,L42  
 L45 12 S L43,L44  
 L46 151 S (C6H16O18P4 OR C6H15O15P3 OR C6H14O12P2)/MF AND 46.150.1/RID  
 L47 9 S 2466-09-3 OR 10380-08-2 OR 13813-62-2 OR 12395-97-0 OR 29444-  
 L48 1 S 7664-38-2  
 L49 10 S L47,L48  
       SEL RN  
 L50 16644 S E68-E77/CRN  
 L51 2775 S L50 AND PMS/CI  
 L52 232 S L51 AND HOMOPOLYMER  
 L53 117 S L52 AND NR>=1  
 L54 115 S L52 NOT L53  
 L55 16 S L54 AND (LI OR H3N OR NA)  
 L56 4 S L55 AND 2/NC  
 L57 8 S L55 AND 3/NC  
 L58 1 S L57 AND K  
 L59 5 S L56,L58



L60 2543 S L51 NOT L52-L59  
 L61 1751 S L60 AND NR>=1  
 L62 792 S L60 NOT L61  
 L63 363 S L62 NOT (C2H4O OR C3H6O)  
 L64 142 S L63 NOT (N OR S OR SI)/ELS  
 L65 2697 S L51 AND H3O4P  
 L66 5 S L65 AND 1/NC  
 L67 2 S L40 NOT L45  
 L68 1 S L67 AND H3O3P  
 L69 22 S L45,L49,L68  
 L70 170 S L46,L69  
 L71 10 S (O7P2 OR O13P4 OR O16P5 OR O19P6 OR O22P7 OR O25P8 OR L28P9 O  
 L72 180 S L70,L71  
 E HEXAMETAPHOSPHATE/CN  
 L73 1 S E3  
 L74 2 S E4,E5  
 E OPOP/ES  
 L75 1511 S (OPOP OR OPOPOP OR OPOPOP OR OPOPOPOP OR OPOPOPOPOP OR OPOPOP  
 L76 177 S L75 AND 1/NR AND 1/NC  
 L77 162 S L76 NOT (ETHOXY OR METHOXY OR PROPOXY)  
 L78 134 S L77 NOT (TIS OR AYS OR MNS)/CI  
 L79 132 S L78 NOT CCS/CI  
 L80 50 S L79 NOT (C OR F OR CL OR S OR N)/ELS  
 L81 41 S L80 NOT RPS/CI  
 L82 35 S L81 NOT ION  
 L83 9 S L82 AND (H2O6P2 OR H3O8P3 OR H4O12P4 OR H5O15P5 OR H6O18P6 OR  
 L84 6 S L83 NOT 3-4/P  
 L85 3 S L83 NOT L84  
 L86 1 S L85 NOT OPOPOP/ES  
 L87 7 S L84,L86  
 L88 2 S L75 AND H9O27P9  
 L89 268 S L75 AND H3O9P3  
 L90 1 S L89 AND 1/NC  
 L91 8 S L87,L90  
 L92 188 S L72,L91  
 E GLASS/CN  
 L93 1 S E56  
 L94 189 S L92,L93

FILE 'HCAPLUS' ENTERED AT 09:07:16 ON 04 JUN 2003

L95 81393 S L94  
 L96 5383 S INOSITOL(L) (DIPHOSPH? OR TRIPHOS? OR TETRAPHOS? OR PENTAPHOS?  
 L97 82822 S IMIDODIPHOS? OR GUANYLIMIDODIPHOS? OR ADENYLYLIMIDODIPHOS? OR  
 L98 111 S (ETIDRONIC OR PAMIDRONIC) ()ACID

FILE 'REGISTRY' ENTERED AT 09:08:54 ON 04 JUN 2003

FILE 'HCAPLUS' ENTERED AT 09:11:24 ON 04 JUN 2003

L99 991 S AMIDRONIC ACID OR (GUANYL? OR ADENYL?) ()IMIDODIPHOSPH?  
 L100 156796 S L95-L98,L99  
 L101 1483 S L24-L33 AND L100  
 L102 17 S L10-L13 AND L100  
 L103 2 S L101 AND L102  
 L104 2 S L38,L103  
 L105 15 S L102 NOT L104  
 L106 91 S L100 (L) THU/RL AND L101  
 L107 66 S L95 (L) THU/RL AND L106  
 L108 12 S L107 AND ?ALZHEIM?  
 L109 9 S L107 AND ?AMYLO?  
 L110 19 S L108,L109  
 L111 1277 S L101 AND (PD<=20000501 OR PRD<=20000501 OR AD<=20000501)  
 L112 63 S L111 AND L106  
 L113 10 S L110 AND L112

SEL DN AN 2 3 4 5 6  
L114 5 S E1-E15 AND L113  
L115 53 S L112 NOT L113  
SEL DN AN 16 28 27 42  
L116 4 S E16-E27 AND L115  
L117 10 S L114,L116,L104  
SEL HIT RN

FILE 'REGISTRY' ENTERED AT 09:25:37 ON 04 JUN 2003  
L118 16 S E28-E43  
L119 15 S L118 NOT UNSPECIFIED

FILE 'REGISTRY' ENTERED AT 09:26:07 ON 04 JUN 2003

FILE 'HCAPLUS' ENTERED AT 09:26:19 ON 04 JUN 2003

FILE 'MEDLINE' ENTERED AT 09:26:56 ON 04 JUN 2003  
L120 12812 S L94  
L121 72485 S L97,L98,L99  
E PYROPHOSPHATE/CT  
E E20+ALL  
E E2+ALL  
L122 4763 S E17,E21  
L123 4763 S E17,E21/CN  
E E16+ALL  
L124 638 S E16  
L125 638 S E16/CN  
L126 79693 S L120-L125  
E ALZHEIMER/CT  
E E8+ALL  
L127 37424 S E12+NT OR E46+NT OR E47+NT OR E48+NT OR E49+NT OR E50+NT OR E  
L128 119 S L126 AND L127  
L129 105 S L128 AND PY<=2000  
L130 4 S L129 NOT AB/FA  
L131 101 S L129 NOT L130  
L132 3 S L131 AND OXIDATIVE STRESS  
L133 98 S L131 NOT L132  
SEL DN AN 29 30 58 65 67 78 80 81 85 86 90  
L134 11 S E1-E33  
L135 11 S L134 AND L120-L134  
E FREYW/AU  
E FREY W/AU  
L136 109 S E3,E4,E12,E13,E21,E22  
E FAWCETT J/AU  
L137 284 S E3,E13  
E E24  
L138 1 S E5  
L139 5 S L126 AND L136-L138  
L140 1 S L139 AND ?ALZHEIM?  
L141 1 S L139 AND L128  
L142 1 S L140,L141  
L143 12 S L135,L142

FILE 'MEDLINE' ENTERED AT 09:45:59 ON 04 JUN 2003

FILE 'BIOSIS' ENTERED AT 09:46:08 ON 04 JUN 2003  
E FREY W/AU  
L144 178 S E3,E7-E12,E20-E24  
E FAWCETT J/AU  
L145 119 S E3,E12  
L146 11 S E28,E30  
L147 295 S L144-L146  
L148 2 S L147 AND L94

FILE 'BIOSIS' ENTERED AT 09:47:31 ON 04 JUN 2003

FILE 'WPIX' ENTERED AT 09:47:56 ON 04 JUN 2003

E FREY W/AU  
L149 120 S E3,E7  
E FAWCETT J/AU  
L150 4 S E3,E7  
E HEALTHPART/PA  
L151 1 S E4,E5  
L152 1 S L149,L150 AND L151  
E A61K031-66/IC,ICM,ICS  
L153 2927 S E3-E20  
E A61K031-706/IC,ICM,ICS  
L154 150 S E3-E7  
L155 319 S E18-E20  
L156 54 S E24-E26  
E A61K031-7105/IC,ICM,ICS  
L157 225 S E3-E5  
L158 13743 S (B05-B01G OR B05-B01J OR B05-B01P OR C05-B01G OR B05-B01J OR  
L159 15870 S L153-L158  
L160 6475 S (?PYROPHOSPH? OR ?PYRO PHOSPH?)/BIX  
L161 22154 S L159,L160  
L162 3 S L149-L151 AND L161  
L163 3 S L152,L162  
L164 389 S L161 AND ?ALZHEIM?/BIX  
L165 383 S L161 AND (B14-J01A4 OR C14-J01A4 OR B12-G04A OR C12-G04A)/MC  
L166 251 S P444/M0,M1,M2,M3,M4,M5,M6 AND L161  
L167 574 S L164-L166  
L168 27 S L167 AND L160  
SEL DN AN 14  
L169 1 S E1-E2  
L170 545 S L167 NOT L163,L168-L169  
L171 223 S L170 AND L153  
L172 57 S L171 AND (ALZHEIM? OR AMYLO?)/TI  
L173 103 S ((B115 OR B415 OR B515 OR B615)(S)(B702 OR B713))/M0,M1,M2,M3  
L174 25 S L172 AND L173  
L175 32 S L172 NOT L174  
L176 78 S L173 NOT L174,L175  
L177 88 S L171 NOT L172-L176  
L178 3 S L163,L169 AND L149-L177

FILE 'WPIX' ENTERED AT 10:18:01 ON 04 JUN 2003

FILE 'DPCI' ENTERED AT 10:18:15 ON 04 JUN 2003  
E WO2001082932/PN